

2020

Cervical Cancer Screening Participation among Women of Russian, Somali, and Kurdish Origin Compared with the General Finnish Population: A Register-Based Study

Idehen, Esther E

MDPI AG

Tieteelliset aikakauslehtiartikkelit

© 2020 by the authors

CC BY <http://creativecommons.org/licenses/by/4.0/>

<http://dx.doi.org/10.3390/ijerph17217899>


<https://erepo.uef.fi/handle/123456789/24194>

Downloaded from University of Eastern Finland's eRepository



Article

Cervical Cancer Screening Participation among Women of Russian, Somali, and Kurdish Origin Compared with the General Finnish Population: A Register-Based Study

Esther E. Idehen ^{1,*} , Anni Virtanen ^{2,3}, Eero Lilja ⁴, Tomi-Pekka Tuomainen ¹, Tellervo Korhonen ⁵ and Päivikki Koponen ⁶

¹ Institute of Public Health and Clinical Nutrition, Faculty of Health Sciences, School of Medicine, University of Eastern Finland, Yliopistonranta 1, P.O. Box 1627, 70211 Kuopio, Finland; tomi-pekka.tuomainen@uef.fi

² Finnish Cancer Registry, Unioninkatu 22, 00130 Helsinki, Finland; anni.virtanen@cancer.fi

³ Department of Pathology, University of Helsinki and HUS Diagnostic Center, Helsinki, University Hospital, Haartmaninkatu 3, 00029 HUS Helsinki, Finland

⁴ Department of Welfare, Finnish Institute for Health and Welfare (THL), P.O. Box 30, FI-00271 Helsinki, Finland; eero.lilja@thl.fi

⁵ Institute for Molecular Medicine Finland (FIMM) University of Helsinki, P.O. Box 20, 00014 Helsinki, Finland; tellervo.korhonen@helsinki.fi

⁶ Department of Public Health Solutions, Finnish Institute for Health and Welfare (THL), P.O. Box 30, FI-00271 Helsinki, Finland; Paivikki.koponen@thl.fi

* Correspondence: estheri@uef.fi; Tel.: +358-503678612

Received: 8 September 2020; Accepted: 24 October 2020; Published: 28 October 2020



Abstract: Migrant-origin women are less prone to cervical screening uptake compared with host populations. This study examined cervical cancer screening participation and factors associated with it in the Finnish mass screening program during 2008–2012 in women of Russian, Somali and Kurdish origin compared with the general Finnish population (Finns) in Finland. The study population consists of samples from the Finnish Migrant Health and Well-being Study 2010–2012 and Health 2011 Survey; aged 30–64 ($n = 2579$). Data from the Finnish screening register linked with other population-based registry data were utilized. For statistical analysis we employed logistic regression. Age-adjusted screening participation rates were Russians 63% (95% CI: 59.9–66.6), Somalis 19% (16.4–21.6), Kurds 69% (66.6–71.1), and Finns 67% (63.3–69.8). In the multiple-adjusted model with Finns as the reference; odds ratios for screening were among Russians 0.92 (0.74–1.16), Somalis 0.16 (0.11–0.22), and Kurds 1.37 (1.02–1.83). Among all women, the substantial factor for increased screening likelihood was hospital care related to pregnancy/birth 1.73 (1.27–2.35), gynecological 2.47 (1.65–3.68), or other reasons 1.53 (1.12–2.08). Screening participation was lower among students and retirees. In conclusion, screening among the migrant-origin women varies, being significantly lowest among Somalis compared with Finns. Efforts using culturally tailored/population-specific approaches may be beneficial in increasing screening participation among women of migrant-origin.

Keywords: cervical cancer screening; cancer prevention; disparities; healthcare service utilization; health inequities; migrant health; women health; public health; population-based study; quantitative research

1. Introduction

Due to screening programs and through regular cervical cytological Papanicolaou (Pap) testing offered to all at-risk women as an effective preventive measure against the onset of cervical cancer [1–4];

incidence and mortality rates from cervical cancer in some higher-income countries have been significantly declining [5,6]. However, the disease's risk is significant in women not adhering to screening recommendations [4,7–9], explicitly found among some migrant-origin populations [4,10–13]. Therefore, a well-organized screening program and active screening participation within the target population are imperative for comprehensive coverage and reducing the disease incidence and mortality rates [1,2,4,7].

Migrant populations have increased globally, including in Finland [14,15], and the health status of these population groups has become a public health concern, highlighting the need to strengthen healthcare systems [16–19]. Studies have revealed the underutilization of cervical cancer screening services among some migrant-origin groups compared with host populations [10–13,20–26]. Consequently, persisting health inequities will increase national healthcare expenditure [27]. Studies have linked some barriers to cervical cancer screening participation at the individual and the screening system levels [28–30].

Some individual barriers to cervical cancer screening participation include low socioeconomic status, being unmarried, and unemployed, illiteracy, and limited language skills [11,21,22,25,30,31]; inadequate understanding or unawareness about screening/purpose, cancer risk in the host country/country of origin, and fear of cancer or screening [30–35]; cultural/religious beliefs; and unpleasant screening experiences, such as pain, and embarrassment owing to Female Genital Mutilation (FGM), and/or obesity [26,36–39]. Some other individual barriers are being an older/nulliparous woman, not using healthcare services related to gynecological, reproductive, or pregnancy [21,30,40–42], and issues related to migration and residential areas [22,25,26,33,43]. System barriers include women's distrust in healthcare authorities and personnel, unavailability of female screeners, inaccessibility to healthcare, and interpretation services [25,32,33,35,39,43].

In Finland, municipalities are obliged to offer free-of-charge cervical cancer screening to eligible women aged 30–60, with a five-year interval [44]. Additionally, some municipalities invite women aged from 25 to 65. Eligible women are identified from the Finnish National Population Registry covering all Finnish citizens and foreign citizens residing in Finland on a permanent or temporary basis (excluding only undocumented migrants), and personal invitations are sent by mail [44]. The invitation coverage is almost 100% among women aged 30–60 [44]. Presently, the overall participation rate in the cervical screening program is about 70% [44], although the aim is to achieve 80–85% screening coverage in the population [44]. In addition to the organized program, opportunistic screening, i.e., the testing of non-symptomatic persons outside the program, is widespread in the country [45]. Opportunistic screening is not included in any national register [46].

Previous studies in Finland demonstrated disparities and lower cervical cancer screening participation among migrant-origin women [20,21] and those with a non-native mother tongue [30,46–48], compared with the general Finnish population (hereafter referred to as Finns). However, register-based studies about cervical cancer screening participation in the mass screening program among various migrant-origin populations are limited. This study examined cervical cancer screening participation and factors associated with it in the Finnish mass screening program during 2008–2012 in women of Russian, Somali, and Kurdish origin compared with the general Finnish population in Finland.

2. Material and Methods

2.1. Study Population

This study population includes women from the samples of the Migrant Health and Well-being Study 2010–2012 (Maamu) [49] and the Health 2011 Survey [50], both of which were carried out by the Finnish Institute for Health and Welfare (THL). The sample from the Maamu study comprises migrant-origin populations of 1998 Russians (1230 women), 1963 Somalis (1020 women), and 1948 Kurdish-origin (819 women) (briefly, hereafter referred to as Kurds). The study stratified random sample was drawn from the Finnish National Population Registry; inclusion criteria were as follows: age 18–64,

residence in one of the six cities (Helsinki, Vantaa, Espoo, Tampere, Turku, and Vaasa) with a high proportion of the migrant-origin population, and residence for at least a year in Finland. The country of birth was Somalia, Russia or the former Soviet Union, and Iran or Iraq. The native language was Russian or Finnish for the Russian group and Kurdish Sorani for the Kurdish-origin group.

The reference population (Finns) consists of women from the Health 2011 Survey sample, including the same age groups and cities as in the Maamu Study. We identified a small portion of women from the study groups who had no information about the invitation to the mass screening; we excluded them from further analysis. This study sample was narrowed to women aged 30–64 and those migrants who arrived in Finland before 2008. Thus, all women included in the analyses were: 816 Russians, 523 Somalis, 451 Kurds, and 789 Finns, who had received at least one invitation from the organized screening program (Figure 1).




Figure 1. Flow chart of the study population. (Six cities: Helsinki, Espoo, Vantaa, Turku, Tampere, and Vaasa).

2.2. Data Sources and Variables

We employed data from the: (1) National Population Register [51], (2) Mass Screening Registry [52], (3) Care Register for Health Care [53], (4) Medical Birth Register [54], (5) Register of Induced Abortions [55], (6) Statistics Finland [56] and, (7) Social insurance institution of Finland (Kela), [57]. The data from different sources for persons in the Maamu and Health 2011 Survey samples were linked using the unique personal identity codes [58] given to Finland's legal residents.

2.3. Study Outcomes

The study outcome measure was participation in the organized cervical cancer screening program among all eligible women invited during 2008–2012 in Finland. In Finland, the invitation to screening is usually valid for a year, after these women are regarded as non-attendees.

2.4. Variable Definitions

Invitation and participation data for the screening program came from the Mass Screening Registry [52]; they were dichotomously coded as yes vs. no; yes indicated a positive response to the latest invitation 2008–2012. We obtained data on age from the population register; our three categories were 30–39, 40–49, and 50–64. Education data came from Statistics Finland and was dichotomously coded as at least high school or equivalent vs. upper secondary or less by 2012.

Marital status data was from the population register on the date the sample was drawn in 2010; the status was dichotomously coded as married or in a civil union vs. any other or unknown. The year of migration to Finland came from the population register being dichotomously coded as 1970–1997 vs. 1998–2007. Employment status data were from Statistics Finland for the invitation year or the closest available year (2009, 2010, or 2011). When unknown, the data were supplemented with information on Kela benefits from 2010–2012. This data was coded into five categories: employed, childcare at home, student, retired, and unemployed/unknown.

Information on the item having moved from one municipality to another in Finland came from the Population Register and was dichotomously coded for 2008–2011 as a change in residence municipality vs. no change during that period. This information was only available for women with a migrant background, who are likely to move more often than the general population. Information on the item having stayed abroad for over one year from Finland was obtained from the Population register and dichotomously coded as having stayed abroad for at least one year vs. less or not at all, between 2009 and 2011.

The number of births (the number of deliveries an individual had) in Finland came from the Medical Birth Register for 1987–2012 and was coded as none, 1 or 2, and 3 or more births. Only those births before the screening invitation year were considered. The number of abortions in Finland was obtained from the Register of Induced Abortions and was dichotomously coded as follows: none vs. one or more, covering 1983–2012. Only abortions before the screening invitation year were considered.

Hospital care in Finland was identified from the Care Register for Health Care as having been in secondary healthcare as inpatient or outpatient: these recorded visits were categorized as: no care; pregnancy or birth-related care (International Classification of Diseases (ICD 10) codes O00–O08, O10–O16, O20–O29, O30–O48, O60–O75, O80–O92, O94–O99); other gynecological care (ICD10: N70–77, N80–N98, and gynecological cancers C51–C58, D06, D07); and hospital care for any other reason. Only hospital care from 1994 until the year before the screening invitation was considered.

2.5. Statistical Analyses

For statistical analyses, we utilized SAS 9.3 and SUDAAN 11.0.3 software [59]. The study population's age-adjusted main descriptive characteristics were explored by country of origin. Furthermore, logistic regression was used, adjusting for age to explore factors associated with participation. The study results are shown as odds ratios (ORs), 95% confidence intervals (CIs), *p*-values, and model-adjusted screening participation percentages. We used the Akaike information criterion (AIC) [60] to determine the best predictive model for screening participation. In the model selection, we considered all the main effects and how these interacted with the country of origin.

For the final model, the selected variables were country of origin, age, education, employment status, staying abroad, hospital care in Finland, and the interaction between the country of origin, and year of migration. The statistical significance was assessed with the Satterthwaite-adjusted *F*-value. Age-adjusted proportions were calculated using predictive margins [61]. In all the analyses, we applied

Finite Population Correction (FPC) to the migrant-origin groups and inverse probability weights (IPWs) [62], due to unequal sampling probabilities within the study groups.

2.6. Ethical Considerations

The coordinating ethics committee of the Hospital District of Helsinki and Uusimaa in Finland approved the use of register data for both the Maamu Study and Health 2011 Survey samples, including the linkage of data from different registers (decisions #325/13/03/00/2009 and 45/13/03/00/11). All data was analyzed and stored at THL, following THL data safety regulations and complying with the European Union General Data Protection Regulation [63].

3. Results

3.1. Characteristics

Table 1 displays an overview of the characteristics of the study populations. Women of Russian origin had a higher proportion of high education level (79%) and employment (60%), compared with Kurds (33% and 32%), Somalis (19% and 26%); the highest percentages were those of the Finns (87% and 81%). The Kurds had the highest proportion of being married or in a civil union (76%), compared to Russians (54%), Somalis (65%), and Finns (48%).

Table 1. Mass screening participation in the study population and baseline characteristics of the women meeting the inclusion criteria (aged 30–64) and the migrant origin groups who have moved to Finland between the years 1970 and 2007) by country of origin: weighted and age-adjusted proportions.

Characteristics (n = 2644)	Russian (n = 837)		Somali (n = 532)		Kurdish (n = 454)		Finns (n = 821)	
	%	95%CI ¹	%	95%CI	%	95%CI	%	95%CI
Mass screening participation								
Variable/description								
No invitation	2.6	1.7–4.1	2.1	1.3–3.4	0.9	0.4–1.8	3.5	2.5–4.9
Invited, not participated	34.0	30.8–37.4	79.0	76.2–81.6	30.2	28.1–32.4	29.9	26.8–33.1
Invited, participated	63.3	59.9–66.6	18.9	16.4–21.6	68.9	66.6–71.1	66.6	63.3–69.8
Age								
30–39	29.1	26.0–32.3	44.0	40.8–47.3	43.5	41.0–45.9	28.7	25.8–31.9
40–49	34.3	31.0–37.6	36.2	33.1–39.4	39.9	37.5–42.4	28.5	25.5–31.7
50–64	36.7	33.4–40.1	19.7	17.3–22.4	16.6	14.9–18.5	42.8	39.4–46.2
Education ²								
High school or higher	78.7	75.7–81.4	18.8	16.4–21.4	33.0	30.7–35.3	87.0	84.7–89.1
Less than high school/unknown	21.3	18.6–24.3	81.2	78.6–83.6	67.0	64.7–69.3	13.0	10.9–15.3
Marital status								
Married or in a civil union	53.7	50.1–57.1	65.4	62.2–68.5	76.5	74.4–78.6	48.5	45.0–51.9
Other/Unknown	46.3	42.9–49.9	34.6	31.5–37.8	23.5	21.4–25.6	51.5	48.1–55.0
Employment status								
Employed	60.1	56.6–63.5	26.0	23.2–29.0	32.1	29.9–34.5	81.1	78.3–83.5
Childcare at home	5.6	4.2–7.4	5.3	4.0–6.9	5.8	4.8–7.1	1.7	1.0–2.9
Student	6.1	4.6–8.2	22.1	19.8–24.5	11.9	10.6–13.4	2.5	1.5–3.9
Retired	3.7	2.6–5.2	6.6	5.0–8.5	13.2	11.5–15.2	5.1	3.9–6.7
Unemployed/unknown	24.5	21.7–27.6	40.0	36.9–43.2	36.9	34.4–39.4	9.6	7.9–11.7
Year of migration								
1970–1997	52.0	48.5–55.4	66.2	63.0–69.2	43.7	41.3–46.2		
1998–2007	48.0	44.6–51.5	33.8	30.8–37.0	56.3	53.8–58.7		
Had moved from one municipality to another								
No	94.6	92.5–96.1	97.0	95.4–98.0	95.8	94.9–96.6		Na ³
Yes	5.4	3.9–7.5	3.0	2.0–4.6	4.2	3.4–5.1		

Table 1. Cont.

Characteristics (n = 2644)	Russian (n = 837)		Somali (n = 532)		Kurdish (n = 454)		Finns (n = 821)	
Had stayed abroad for over one-year ⁴								
No	97.4	96.0–98.3	92.3	90.2–94.0	93.1	91.5–94.4	97.7	96.4–98.6
Yes	2.6	1.7–4.0	7.7	6.0–9.8	6.9	5.6–8.5	2.3	1.4–3.6
Number of births given in Finland								
None	56.3	53.1–59.5	36.8	34.0–39.7	42.2	40–44.5	42.4	39.2–45.8
1–2	40.8	37.6–44.0	15.5	13.4–18.0	47.3	44.9–49.6	45.3	41.9–48.8
3 or more	2.9	2.0–4.4	47.7	44.7–50.7	10.5	9.3–11.8	12.2	10.1–14.8
Number of abortions in Finland								
None	81.1	78.2–83.7	90.3	88.3–92.0	77.2	75.1–79.2	82.1	79.2–84.6
1 or more	18.9	16.3–21.8	9.7	8.0–11.7	22.8	20.8–24.9	17.9	15.4–20.8
Hospital care in Finland ⁵								
None	20.1	17.4–23.0	4.7	3.4–6.5	4.6	3.5–6.1	14.8	12.5–17.3
Pregnancy/birth-related care	41.8	38.6–45.0	61.3	58.4–64.2	59.3	57.0–61.6	38.6	35.4–41.9
Other gynecological reason	10.2	8.3–12.5	10.0	8.1–12.4	9.4	8.0–11.0	11.7	9.7–14.0
Other reason	28.0	25.1–31.1	23.9	21.4–26.7	26.7	24.4–29.1	35.0	31.9–38.2

¹ 95% CI = 95% confidence interval; ² High school education registered in Finland; ³ Na = not available; ⁴ Had stayed abroad for a period of over one year between the years 2008–2011; ⁵ Hospital care in Finland = inpatient or outpatient care in secondary or tertiary health care for the specified reasons.

3.2. Screening Participation in the Age-Adjusted Analysis

The age-adjusted screening participation rate among women invited to the screening was highest amongst the Kurds (69%), followed by Finns (67%) and Russians (63%); it was markedly lowest among Somalis (19%).

Among the Russians, a higher education level (OR = 1.52; 95% confidence interval (CI): 1.06–2.17), and having had hospital care related to pregnancy/birth (OR = 1.97; 95% CI: 1.29–3.02), or for other gynecological reasons (OR = 2.09; 95% CI: 1.13–3.86) compared to no history of hospital care in Finland were associated with screening participation. Additionally, increased participation with advanced age was observed. Retirees participated less (OR = 0.39; 95% CI: 0.17–0.87) compared with employed women.

Among the Somalis, higher screening participation was associated with a recent migration to Finland (OR = 1.56; 95% CI: 1.10–2.21), and those aged 40–49 (OR = 1.49; 95% CI: 1.03–2.16); age 30–39 as the reference. Participation was lower among students (OR = 0.37; 95% CI: 0.22–0.62), with employed women as the reference. Participation decreased as the number of births increased (OR = 0.57; 95% CI: 0.36–0.92).

Among the Kurds, factors associated with higher screening participation were age 40–49 (OR = 1.64; 95% CI: 1.30–2.06) as well as hospital care relating to pregnancy/birth (OR = 3.54; 95% CI: 1.92–6.53), gynecological (OR = 3.01; 95% CI: 1.49–6.08), or for other reasons (OR = 2.70; 95% CI: 1.42–5.13), compared to no history of hospital care in Finland. Other factors were having lived in one municipality in Finland compared to having moved from one municipality to another (OR = 1.79; 95% CI: 1.16–2.75); being married or in a civil union (OR = 1.57; 95% CI: 1.24–1.99), and higher education level (OR = 1.41; 95% CI: 1.13–1.76). Lower participation was associated with being a student (OR = 0.68; 95% CI: 0.48–0.95), retired (OR = 0.54; 95% CI: 0.37–0.78), or unemployed/unknown (OR = 0.66; 95% CI: 0.51–0.86).

Among the Finns, older age (50–64), (OR = 1.96; 95% CI: 1.35–2.83), higher education level (OR = 1.87; 95% CI: 1.20–2.89), and hospital care due to other gynecological reasons (OR = 2.77; 95% CI: 1.40–5.49), as well as staying abroad for less than a year (OR = 4.86; 95% CI: 1.62–14.57) compared to staying abroad for at least a year during the study period were associated with higher screening participation (Table 2).

Table 2. Age-adjusted participation in mass cervical cancer screening by country of origin (includes only women invited to mass screening).

Total Women (n = 2579)	Russian (n = 816)			Somali (n = 523)			Kurdish (n = 451)			Finns (n = 789)		
Mass screening participation	%	95% CI ¹		%	95% CI		%	95% CI		%	95% CI	
Invited, participated	63.3	59.9–66.6		18.9	16.4–21.6		68.9	66.6–71.1		66.6	63.3–69.8	
Variable/description	%	OR ² (95%CI)	p	%	OR (95%CI)	p	%	OR (95%CI)	p	%	OR (95%CI)	p
Age												
30–39	60.1	1.00		16.8	1.00		63.9	1.00		62.2	1.00	
40–49	67.0	1.35 (0.93–1.96)	0.120	23.1	1.49 (1.03–2.16)	0.027	74.4	1.64 (1.30–2.06)	<0.001	67.5	1.26 (0.86–1.85)	0.001
50–64	68.4	1.44 (1.00–2.07)		14.7	0.85 (0.53–1.37)		67.5	1.17 (0.89–1.54)		76.3	1.96 (1.35–2.83)	
Education ³												
Upper secondary/less	57.9	1.00		19.4	1.00		66.1	1.00		57.4	1.00	
High school or more	67.6	1.52 (1.06–2.17)	0.023	15.9	0.78 (0.50–1.24)	0.300	73.3	1.41 (1.13–1.76)	0.002	71.3	1.87 (1.20–2.89)	0.005
Marital status												
Other/unknown	64.6	1.00		19.3	1.00		61.0	1.00		69.3	1.00	
Married/in a civil union	66.3	1.08 (0.80–1.46)	0.612	18.4	0.94 (0.66–1.34)	0.736	70.9	1.57 (1.24–1.99)	<0.001	69.9	1.03 (0.76–1.40)	0.848
Employment status												
Employed	66.9	1.00		22.7	1.00		74.0	1.00		70.8	1.00	
Childcare at home	62.0	0.80 (0.42–1.54)		27.4	1.28 (0.64–2.56)		75.5	1.08 (0.7–1.68)		71.3	1.02 (0.29–3.63)	
Student	65.7	0.95 (0.48–1.88)	0.195	9.9	0.37 (0.22–0.62)	0.001	65.9	0.68 (0.48–0.95)	0.001	51.1	0.42 (0.15–1.18)	0.399
Retired	44.1	0.39 (0.17–0.87)		23.6	1.05 (0.48–2.31)		60.6	0.54 (0.37–0.78)		66.2	0.81 (0.42–1.56)	
Unemployed/unknown	66.0	0.96 (0.67–1.37)		20.7	0.89 (0.56–1.39)		65.4	0.66 (0.51–0.86)		64.3	0.74 (0.43–1.26)	
Year of migration												
1970–1997	68.0	1.00		16.3	1.00		67.4	1.00				
1998–2007	62.6	0.79 (0.58–1.07)	0.129	23.2	1.56 (1.10–2.21)	0.013	69.6	1.11 (0.89–1.38)	0.358			
Had moved from one municipality to another												
Yes	60.0	1.00		26.2	1.00		55.9	1.00		Na ⁴		
No	65.8	1.28 (0.63–2.63)	0.495	18.5	0.64 (0.24–1.72)	0.371	69.3	1.79 (1.16–2.75)	0.008			
Had stayed abroad for over one-year ⁵												
Yes	45.2	1.00		15.8	1.00		60.9	1.00		33.4	1.00	
No	66.0	2.36 (0.91–6.09)	0.076	18.9	1.24 (0.60–2.57)	0.560	69.3	1.46 (0.93–2.27)	0.097	70.4	4.86 (1.62–14.57)	0.005

Table 2. Cont.

Total Women (n = 2579)	Russian (n = 816)			Somali (n = 523)			Kurdish (n = 451)			Finns (n = 789)		
Number of births given in Finland												
None	63.6	1.00		24.2	1.00		68.7	1.00		70.0	1.00	
1–2 births	68.9	1.27 (0.88–1.83)	0.249	20.6	0.81 (0.46–1.42)	0.045	69.9	1.04 (0.80–1.35)	0.342	69.2	0.96 (0.68–1.36)	0.974
3 births or more	56.9	0.75 (0.32–1.76)		15.6	0.57 (0.36–0.92)		64.7	0.83 (0.59–1.17)		69.2	0.96 (0.57–1.6)	
Number of abortions in Finland												
None	64.9	1.00		18.9	1.00		68.4	1.00		70.1	1.00	
1 or more	68.3	1.17 (0.78–1.75)	0.455	17.2	0.90 (0.47–1.70)	0.736	69.6	1.06 (0.84–1.33)	0.636	67.2	0.87 (0.59–1.29)	0.491
Hospital care in Finland ⁶												
None	55.0	1.00		14.9	1.00		41.5	1.00		62.4	1.00	
Pregnancy/birth-related care	70.5	1.97 (1.29–3.02)	0.010	16.7	1.15 (0.50–2.66)	0.061	71.1	3.54 (1.92–6.53)	<0.001	66.8	1.21 (0.73–2.02)	0.025
Other gynecological reason	71.7	2.09 (1.13–3.86)		27.9	2.23 (0.88–5.70)		67.7	3.01 (1.49–6.08)		82.0	2.77 (1.40–5.49)	
Other reason	62.7	1.38 (0.89–2.14)		22.9	1.71 (0.69–4.24)		65.3	2.70 (1.42–5.13)		70.5	1.45 (0.87–2.41)	

¹ 95% CI = 95% confidence interval; ² OR = odds ratio; ³ high school education registered in Finland; ⁴ Na = not available; ⁵ had stayed abroad between the years 2008–201; ⁶ Hospital care in Finland = inpatient or outpatient care in secondary or tertiary health care for the specified reasons.

3.3. Screening Participation in the Multiple-Adjusted Analysis

Table 3 provides the results of the multiple-adjusted analysis, with all three migrant-origin groups and Finns combined. When adjusted for age, education, employment status, staying abroad, and hospital care in Finland, screening participation was highest among the Kurds (OR = 1.37; 95% CI: 1.02–1.83) and lowest among the Somalis (OR = 0.16; 95% CI: 0.11–0.22), with Finns as the reference. Participation rates among Russians were similar to those of the Finns (OR = 0.92; 95% CI: 0.74–1.16).

Among all the groups, the most substantial factor associated with increased screening likelihood was hospital care (related to pregnancy or birth, or other gynecological reasons, and other reasons). Other factors were older age, higher education level, and having not stayed abroad for over a year. Concerning employment status, participation was lower among students and retirees compared with employed women.

Table 3. Multiple adjusted model ¹ for participation in mass cervical cancer screening.

Total <i>n</i> of Women (<i>n</i> = 2579)		
Variable/Description	OR (95% CI) ²	<i>p</i>
Study groups		
Finns	1.00	<0.001
Russian	0.92 (0.74–1.16)	
Somali	0.16 (0.11–0.22)	
Kurdish	1.37 (1.02–1.83)	
Age		
30–39	1.00	0.003
40–49	1.30 (1.04–1.62)	
50–64	1.56 (1.20–2.02)	
Education ³		
Upper secondary or less	1.00	0.006
High school or more	1.37 (1.10–1.70)	
Employment status		
Employed	1.00	0.006
Childcare at home	0.95 (0.61–1.48)	
Student	0.59 (0.43–0.83)	
Retired	0.60 (0.40–0.89)	
Unemployed/unknown	0.82 (0.65–1.03)	
Year of migration		
1970–1997	1.00	0.090
1998–2007 among Russian	0.91 (0.66–1.26)	
1998–2007 among Somali	1.69 (1.08–2.65)	
1998–2007 among Kurdish	1.21 (0.82–1.77)	
Had stayed abroad for over one year		
Yes	1.00	0.008
No	1.87 (1.18–2.96)	
Hospital care in Finland ⁴		
None	1.00	<0.001
Pregnancy and birth-related care	1.73 (1.27–2.35)	
Any other gynecological reason	2.47 (1.65–3.68)	
Any other reason	1.53 (1.12–2.08)	

¹ Including study group, age, education, living abroad, hospital care in Finland and employment status, and using the year of migration separately for each migrant group; ² OR = odds ratio, 95% CI = 95% confidence interval; ³ high school education registered in Finland; ⁴ Hospital care in Finland = inpatient or outpatient care in secondary or tertiary health care for the specified reasons.

4. Discussion

This population register-based study examined cervical cancer screening participation and factors associated with it in the Finnish mass screening program during 2008–2012 in women of Russian, Somali, and Kurdish origin compared with the general Finnish population (Finns) in Finland.

Our study revealed differences in the screening uptake amongst migrant-origin women, consistent with previous studies [10–13,22–26,30,46,64]. Our study is the first population register-based study to report participation in the Finnish mass screening program among these specific migrant-origin groups compared with the general Finnish population in Finland to the best of our knowledge. These findings contribute to existing knowledge and enhance some understanding of cervical cancer screening participation among migrant-origin populations. Sequentially, this can guide policymakers in developing cervical cancer screening protocols to improve participation among migrant-origin populations.

In our study, the participation rate was highest among the Kurds, Russians participated similarly to the Finns, and the rate was clearly lowest among the Somalis, even after adjusting for age, education, employment status, staying abroad, and hospital care in Finland. The low cervical cancer screening participation rate observed amongst Somalis accords with earlier results based on self-reported Pap-smear uptake in Finland [20,21]. The results among the Kurds and Russians differed from those of the previous studies. The relatively good Finnish/Swedish (official Finnish languages) skills among the Russians and Kurds compared with Somalis can partly explain these disparities, as shown in our previous self-reports of population-based study based among these groups [21], and a qualitative study among other migrant groups in Finland [30].

In Finland, healthcare services like screening are offered free-of-charge (except with a nominal fee in few cases) “to all legal inhabitants and their families, irrespective of their cultural background and socioeconomic conditions” [65]. Nevertheless, migrants are a heterogeneous population, and their socioeconomic status can vary depending on their occupation, educational level, and reason for migration. The differences in healthcare systems, social structures, and cultures in the migrants’ new environment compared to their previous societies may impact their utilization of healthcare services such as screening [66,67]. In our study, the migrants’ countries of origin and the purpose of their migration to Finland also differ in several ways [49]. A more extended stay in the immigrants’ host country might enable them to get acquainted with the healthcare system. The health care system in Russia may be more similar compared to the Finnish system than the health care services available in Somalia and Iran, or Iraq.

The Somalis’ markedly low screening rate is consistent with those in international qualitative studies [35,38,39]. These studies demonstrated possible barriers hindering screening participation such as childcare, cultural or religious beliefs, fear, inadequate understanding of screening, limited language skills, absence of female healthcare personnel, mistrust in healthcare authorities, unpleasant screening experiences such as pain, and embarrassment due to FGM practices [35,38,39]. Nonetheless, in our study, such variables were missing, as our main goal was to compare the migrant-origin women with the Finns, using register data for participation in the organized cervical cancer screening in Finland.

Generally, both risk and incidence of cervical cancer are significant in women not adhering to screening recommendations [4,7–9], explicitly found among some migrant-origin populations [4,10–13]. The incidence and mortality of cervical cancer are low in Finland [44], but current overall incidence rates might camouflage the risk of cancer in certain subgroups of the increasingly heterogenic population. The low screening participation among the Somali-origin women is a cause for concern when considering the disease’s risk in the country of origin [68]. There is a strong need for analyses of cancer incidence stratified by immigrant groups.

The increased screening participation rate observed among older women agrees with other studies [37,46–48,69,70]; contradictory reports also exist [11,24]. Higher screening participation with increasing age might stem from a better understanding of the importance of cancer screening among older women [37] or the fact that they have received several screening invitations and are thus more familiar with the concept. On the other hand, overall, lower screening participation among younger women has been connected with higher uptake in opportunistic testing [46]. Younger women may skip the invitation to organized screening if they have recently been tested during health care visits connected to sex counselling and contraception [71]. The higher screening participation observed among younger

Kurds compared with the Finns is consistent with previous studies [46–48]. Younger women in the Kurdish group use less often contraceptive methods requiring medical follow-up than younger women in the general population [71]; they are likely less often offered opportunistic screening.

The association between higher education and higher screening participation is in line with previous national findings from Finland [46,47] and self-reported screening test uptake in Finland, here mainly observed among the Kurds [21], and with other international studies [11,24,37]. Higher education and health literacy may thus be associated [72,73]. Additionally, when we used employees as a reference, no differences between the employed and women taking care of children at home emerged, except for students and retirees, who exhibited lower screening participation. In Finland, people can receive a retirement pension at around age 65 [74]. So, the reasons for retirement among the women studied include mainly severe long-term illness or disability, which can influence screening participation. The association of being married or in a civil union with higher screening participation is consistent with previous findings [11,25,47].

The higher likelihood of screening participation observed among all the groups associated with healthcare services utilization related to gynecological reason, pregnancies, or births history agrees with other studies [21,40–42]. Contact with healthcare might enable women to be more acquainted with the healthcare system and obtain disease prevention-related information, thus facilitating healthcare services use such as screening participation [43]. Finally, the association of absence from Finland for an extended period with decreased screening participation is apparent as the screening invitation is sent as a personal invitation letter, which is usually valid for a year [44].

Limitations and Strengths

This study has some limitations. Firstly, it has examined cervical cancer screening participation in the Finnish screening program among only these groups and specific cities in Finland; we cannot generalize these results to all migrant-origin women populations. Secondly, the study used data based on registered information on women invited to the screening within a specific period. We are uncertain whether all the women invited received these mailed invitations; the population register may contain inaccurate addresses. Similarly, missing items in the register data may have caused some bias; unknown education or employment status is an example. Thirdly, this study focused on organized screening uptake but information on the use of opportunistic testing would have provided valuable background information as an explanatory factor to non-participation in the organized program.

This study's key strengths are that it utilized national registries, all of which have proven data quality and coverage in Finland [75–77]. Self-reported screening uptake may be vulnerable to recall bias [78]; in contrast, register-based data can provide more reliable information on screening in the organized program. Furthermore, population-based random survey samples were used to identify women in the three groups, which are among the largest migrant-origin populations in Finland, and the cities selected covered a significant share of these populations. The size of the city-specific sub-samples was determined according to the size of the foreign-born population living in a particular city [49].

5. Conclusions

Despite the free-of-charge mass cervical cancer screening organized in Finland, our study demonstrated that screening participation among the migrant origin women varies, being significantly lowest among Somalis compared with women in the general Finnish population (Finns). Efforts using culturally tailored/population-specific approaches may be beneficial in increasing screening participation among women of migrant origin. More attention is needed to raise awareness about the importance of screening and preventive care in certain groups. These groups comprise women of migrant backgrounds, low socioeconomic status, younger women, students, retirees, and not using healthcare services related to gynecological or/and reproductive reasons. The markedly low participation rate observed among Somali-origin women is a cause for concern and requires further attention. More broadly, qualitative research is needed to explore this group's perceptions regarding

the screening and factors related to social relationships, religion, or culture, which cannot be studied with register-based data.

Author Contributions: Conceptualization, E.E.I., A.V., T.K. and P.K.; data curation, E.L.; formal analysis, E.L.; funding acquisition, E.E.I.; methodology, E.E.I., A.V., E.L. and P.K.; project administration, P.K.; software, E.L.; supervision, T.-P.T. and T.K.; validation, E.E.I., A.V., T.-P.T., T.K. and P.K.; visualization, E.E.I., A.V., T.-P.T., T.K. and P.K.; writing—original draft, E.E.I.; writing—review and editing, E.E.I., A.V., E.L., T.-P.T., T.K. and P.K. All authors have read and agreed to the published version of the manuscript.

Funding: This work was funded by the Juho Vainio Foundation (Grant No. 201910465/202010184) and Cancer Society of North Savo (Grant No. 201700009/201810012), Finland, for personal Ph.D. research working grants (E.E.I.).

Acknowledgments: We warmly acknowledged all the study group experts in the Finnish Institute for Health and Welfare (THL) and Finnish Cancer Registry for their involvement in this study. We acknowledge Gerald G Netto at the University of Eastern Finland for checking the English language of this paper.

Conflicts of Interest: The authors declare no conflict of interest. The funders had no role in the design of this study; in the collection, analyses, or interpretation of data; in the writing of the manuscript; or in the decision to publish the results.

Research Data: Due to data protection regulations, the register data are not openly shared. However, data can be obtained for research purposes. Guidelines for data use are available on the survey website: <https://thl.fi/en/web/thlfi-en/research-and-expertwork/population-studies/migrant-health-and-wellbeing-study-maamu/information-for-researchers>.

References

1. WHO. Screening for Cervical Cancer. Available online: <http://www.who.int/reproductivehealth/topics/cancers/fight-cervical-cancer/en/> (accessed on 6 June 2020).
2. Landy, R.; Pesola, F.; Castañón, A.; Sasieni, P. Impact of cervical screening on cervical cancer mortality: Estimation using stage-specific results from a nested case–control study. *Br. J. Cancer* **2016**, *115*, 1140–1146. [[CrossRef](#)] [[PubMed](#)]
3. Vaccarella, S.; Franceschi, S.; Zaridze, D.; Poljak, M.; Veerus, P.; Plummer, M.; Bray, F. Preventable fractions of cervical cancer via effective screening in six Baltic, central, and eastern European countries 2017–40: A population-based study. *Lancet Oncol.* **2016**, *17*, 1445–1452. [[CrossRef](#)]
4. Spence, A.R.; Alobaid, A.; Drouin, P.; Goggin, P.; Gilbert, L.; Provencher, D.; Tousignant, P.; Hanley, J.A.; Franco, E.L. Screening histories and contact with physicians as determinants of cervical cancer risk in Montreal, Quebec. *Curr. Oncol.* **2014**, *21*, 294–304. [[CrossRef](#)] [[PubMed](#)]
5. WHO. Cervical Cancer. Available online: https://www.who.int/health-topics/cervical-cancer#tab=tab_1 (accessed on 23 May 2020).
6. Bray, F.; Ferlay, J.; Soerjomataram, I.; Siegel, R.L.; Torre, L.A.; Jemal, A. Global cancer statistics 2018: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA Cancer J. Clin.* **2018**, *68*, 394–424. [[CrossRef](#)] [[PubMed](#)]
7. Altobelli, E.; Lattanzi, A. Cervical carcinoma in the European Union: An update on disease burden, screening program state of activation, and coverage as of March 2014. *Int. J. Gynecol. Cancer* **2015**, *25*, 474–483. [[CrossRef](#)]
8. Lonnberg, S.; Nieminen, P.; Luostarinen, T.; Anttila, A. Mortality audit of the Finnish cervical cancer screening program. *Int. J. Cancer* **2013**, *132*, 2134–2140. [[CrossRef](#)]
9. Dugue, P.; Lynge, E.; Rebolj, M. Mortality of non-participants in cervical screening: Register-based cohort study. *Int. J. Cancer* **2014**, *134*, 2674–2682. [[CrossRef](#)]
10. Visioli, C.B.; Crocetti, E.; Zappa, M.; Iossa, A.; Andersson, K.L.; Bulgaresi, P.; Alfieri, A.; Amunni, G. Participation and Risk of High Grade Cytological Lesions Among Immigrants and Italian-Born Women in an Organized Cervical Cancer Screening Program in Central Italy. *J. Immigr. Minor. Health* **2015**, *17*, 670–678. [[CrossRef](#)]
11. Gallo, F.; Caprioglio, A.; Castagno, R.; Ronco, G.; Segnan, N.; Giordano, L. Inequalities in cervical cancer screening utilisation and results: A comparison between Italian natives and immigrants from disadvantaged countries. *Health Policy* **2017**, *121*, 1072–1078. [[CrossRef](#)]
12. Azerkan, F.; Sparen, P.; Sandin, S.; Tillgren, P.; Faxelid, E.; Zendejdel, K. Cervical screening participation and risk among Swedish-born and immigrant women in Sweden. *Int. J. Cancer* **2012**, *30*, 937–947. [[CrossRef](#)]

13. Van Leeuwen, A.W.; de Nooijer, P.; Hop, W.C. Screening for cervical carcinoma: Participation and results for ethnic groups and socioeconomic status. *Cancer Cytopathol.* **2005**, *105*, 270–276. [CrossRef]
14. International Organization for Migration (IOM). World Migration Report 2018. Available online: https://publications.iom.int/system/files/pdf/wmr_2018_en.pdf (accessed on 12 June 2018).
15. Statistics Finland. Numbers and Shares of Persons with Immigrant Background by Area 1990–2017. Available online: http://pxnet2.stat.fi/PXWeb/pxweb/en/Maahanmuuttajat_ja_kotoutuminen/Maahanmuuttajat_ja_kotoutuminen__Maahanmuuttajat_ja_kotoutuminen/007_ulkom_osuudet.px/ (accessed on 15 August 2019).
16. WHO. How Health Systems Can Address Health Inequities Linked to Migration and Ethnicity. Available online: https://www.euro.who.int/data/assets/pdf_file/0005/127526/e94497.pdf (accessed on 12 March 2017).
17. Simon, J.; Kiss, N.; Łaszewska, A.; Mayer, S. *Public Health Aspects of Migrant Health: A Review of the Evidence on Health Status for Labour Migrants in the European Region*; World Health Organisation Regional Office for Europe: København, Denmark, 2015. Available online: http://www.euro.who.int/_data/assets/pdf_file/0003/289245/WHO-HENReport-A5-1-Labour-rev1.pdf?ua=1.2015 (accessed on 1 May 2018).
18. Rechel, B.; Mladovsky, P.; Ingleby, D.; Mackenbach, J.P.; McKee, M. Migration and health in an increasingly diverse Europe. *Lancet* **2013**, *381*, 1235–1245. [CrossRef]
19. Puchner, K.; Karamagioli, E.; Pikouli, A.; Tsiamis, C.; Kalogeropoulos, A.; Kakalou, E.; Pavlidou, E.; Pikoulis, E. Time to rethink refugee and migrant health in Europe: Moving from emergency response to integrated and individualized health care provision for migrants and refugees. *Int. J. Environ. Res. Public Health* **2018**, *15*, 1100. [CrossRef] [PubMed]
20. Idehen, E.E.; Koponen, P.; Härkänen, T.; Kangasniemi, M.; Pietilä, A.-M.; Korhonen, T. Disparities in cervical screening participation: A comparison of Russian, Somali and Kurdish immigrants with the general Finnish population. *Int. J. Equity Health* **2018**, *17*, 56. [CrossRef]
21. Idehen, E.E.; Korhonen, T.; Castaneda, A.; Juntunen, T.; Kangasniemi, M.; Pietilä, A.-M.; Koponen, P. Factors associated with cervical cancer screening participation among immigrants of Russian, Somali and Kurdish origin: A population-based study in Finland. *BMC Women's Health* **2017**, *17*, 19. [CrossRef] [PubMed]
22. Hertzum-Larsen, R.; Kjær, S.K.; Frederiksen, K.; Thomsen, L.T. Participation in cervical cancer screening among immigrants and Danish-born women in Denmark. *Prev. Med.* **2019**, *123*, 55–64. [CrossRef]
23. Brzoska, P.; Aksakal, T.; Yilmaz-Aslan, Y. Utilization of cervical cancer screening among migrants and non-migrants in Germany: Results from a large-scale population survey. *BMC Public Health* **2020**, *20*, 5. [CrossRef] [PubMed]
24. Marlow, L.A.V.; Wardle, J.; Walle, J. Understanding cervical screening non-attendance among ethnic minority women in England. *Br. J. Cancer* **2015**, *113*, 833–839. [CrossRef] [PubMed]
25. Leinonen, M.K.; Campbell, S.; Ursin, G.; Tropé, A.; Nygård, M. Barriers to cervical cancer screening faced by immigrants: A registry-based study of 1.4 million women in Norway. *Eur. J. Public Health* **2017**, *27*, 873–879. [CrossRef] [PubMed]
26. Crampe-Casnabet, C.; Franck, J.; Ringa, V.; Coeuret-Pellicer, M.; Chauvin, P.; Menvielle, G. Role of obesity in differences in cervical cancer screening rates by migration history. The CONSTANCES survey. *Cancer Epidemiol.* **2019**, *58*, 98–103. [CrossRef] [PubMed]
27. Downs, L.S.; Smith, J.S.; Scarinci, I.; Flowers, L.; Parham, G. The disparity of cervical cancer in diverse populations. *Gynecol. Oncol.* **2008**, *109*, S22–S30. [CrossRef]
28. Leinonen, M.K.; Campbell, S.; Klungsoyr, O.; Lönnberg, S.; Hansen, B.T.; Nygård, M. Personal and provider level factors influence participation to cervical cancer screening: A retrospective register-based study of 1.3 million women in Norway. *Prev. Med.* **2017**, *94*, 31–39. [CrossRef] [PubMed]
29. Olesen, S.C.; Butterworth, P.; Jacomb, P.; Tait, R.J. Personal factors influence use of cervical cancer screening services: Epidemiological survey and linked administrative data address the limitations of previous research. *Health Serv. Res.* **2012**, *12*, 34. [CrossRef] [PubMed]
30. Idehen, E.; Pietilä, A.-M.; Kangasniemi, M. Barriers and Facilitators to Cervical Screening among Migrant Women of African Origin: A Qualitative Study in Finland. *Int. J. Environ. Res. Public Health* **2020**, *17*, 7473. [CrossRef] [PubMed]

31. Hulme, J.; Moravac, C.; Ahmad, F.; Cleverly, S.; Lofters, A.; Ginsburg, O.; Dunn, S. "I want to save my life": Conceptions of cervical and breast cancer screening among urban immigrant women of South Asian and Chinese origin. *BMC Public Health* **2016**, *16*, 1–12. [[CrossRef](#)] [[PubMed](#)]
32. Patel, H.; Sherman, S.M.; Tincello, D.; Moss, E.L. Awareness of and attitudes towards cervical cancer prevention among migrant Eastern European women in England. *J. Med. Screen.* **2020**, *27*, 40–47. [[CrossRef](#)]
33. Adunlin, G.; Cyrus, J.W.; Asare, M.; Sabik, L.M. Barriers and facilitators to breast and cervical cancer screening among immigrants in the United States. *J. Immigr. Minor. Health* **2019**, *21*, 606–658. [[CrossRef](#)]
34. Poudel, K.; Sumi, N. Analyzing Awareness on Risk Factors, Barriers and Prevention of Cervical Cancer among Pairs of Nepali High School Students and Their Mothers. *Int. J. Environ. Res. Public Health* **2019**, *16*, 4382. [[CrossRef](#)]
35. Gele, A.A.; Qureshi, S.A.; Kour, P.; Kumar, B.; Diaz, E. Barriers and facilitators to cervical cancer screening among Pakistani and Somali immigrant women in Oslo: A qualitative study. *Int. J. Women's Health* **2017**, *9*, 487–496. [[CrossRef](#)]
36. Marlow, L.; McBride, E.; Varnes, L.; Waller, J. Barriers to cervical screening among older women from hard-to-reach groups: A qualitative study in England. *BMC Women's Health* **2019**, *19*, 38. [[CrossRef](#)]
37. Ekechi, C.; Olaitan, A.; Ellis, R.; Koris, J.; Amajuoyi, A.; Marlow, L.A.V. Knowledge of cervical cancer and attendance at cervical cancer screening: A survey of Black women in London. *BMC Public Health* **2014**, *14*, 1096. [[CrossRef](#)] [[PubMed](#)]
38. Addawe, M.A.; Mburu, C.; Madar, A.A. Barriers to cervical cancer screening: A qualitative study among Somali women in Oslo Norway. *Health Prim. Care* **2018**, *2*, 1–5. [[CrossRef](#)]
39. Salad, J.; Verdonk, P.; de Boer, F.; Abma, T.A. "A Somali girl is Muslim and does not have premarital sex. Is vaccination really necessary?" A qualitative study into the perceptions of Somali women in the Netherlands about the prevention of cervical cancer. *Int. J. Equity Health* **2015**, *14*, 68. [[CrossRef](#)] [[PubMed](#)]
40. Greene, M.Z.; Hughes, T.L.; Sommers, M.S.; Hanlon, A.; Meghani, S.H. Association of pregnancy history and cervical cancer screening in a community sample of sexual minority women. *J. Women's Health* **2019**, *28*, 526–534. [[CrossRef](#)]
41. Grange, G.; Malvy, D.; Lancon, F.; Gaudin, A.; El Hasnaoui, A. Factors associated with regular cervical cancer screening. *Int. J. Gynaecol. Obstet.* **2008**, *102*, 28–33. [[CrossRef](#)]
42. Cerigo, H.; Coutlee, F.; Franco, E.L.; Brassard, P. Factors associated with cervical cancer screening uptake among Inuit women in Nunavik, Quebec, Canada. *BMC Public Health* **2013**, *13*, 438. [[CrossRef](#)]
43. Lofters, A.K.; Moineddin, R.; Hwang, S.W.; Glazier, R.H. Predictors of low cervical cancer screening among immigrant women in Ontario, Canada. *BMC Women's Health* **2011**, *11*, 20. [[CrossRef](#)]
44. Finnish Cancer Registry. Screening Statistics. 2020. Available online: <https://cancerregistry.fi/statistics/screening-statistics/> (accessed on 19 October 2020).
45. Salo, H.; Nieminen, P.; Kilpi, T.; Auranen, K.; Leino, T.; Vänskä, S.; Tiihonen, P.; Lehtinen, M.; Anttila, A. Divergent coverage, frequency and costs of organised and opportunistic Pap testing in Finland. *Int. J. Cancer* **2014**, *135*, 204–213. [[CrossRef](#)]
46. Pankakoski, M.; Heinävaara, S.; Anttila, A.; Sarkeala, T. Differences in cervical test coverage by age, socioeconomic status, ethnic origin and municipality type—A nationwide register-based study. *Prev. Med.* **2020**, *139*, 106219. [[CrossRef](#)]
47. Virtanen, A.; Anttila, A.; Luostarinen, T.; Malila, N.; Nieminen, P. Improving cervical cancer screening attendance in Finland. *Int. J. Cancer* **2015**, *118*, E677–E684. [[CrossRef](#)]
48. Virtanen, A.; Anttila, A.; Luostarinen, T.; Nieminen, P. Self-sampling versus reminder letter: Effects on cervical cancer screening attendance and coverage in Finland. *Int. J. Cancer* **2011**, *128*, 2681–2687. [[CrossRef](#)]
49. Castaneda, A.; Rask, S.; Härkänen, T.; Juntunen, T.; Skogberg, N.; Mölsä, M.; Tolonen, H.; Koskinen, S.; Koponen, P. Enhancing Survey Participation among Foreign-Born Populations. *Finn. Yearb. Popul. Res.* **2018**, *53*, 89–110. Available online: <https://journal.fi/fypr> (accessed on 5 June 2019). [[CrossRef](#)]
50. Lundqvist, A.; Mäki-Opas, T. Health 2011 Survey—Methods. National Institute for Health and Welfare, Report 58-6. 2016. Available online: <http://urn.fi/URN:ISBN:978-952-302-669-8,8/2016> (accessed on 5 June 2019).
51. Population Finland. Population Information System. Available online: <https://vrk.fi/en/population-information-system> (accessed on 5 June 2019).
52. Finnish Cancer Registry. Mass Screening—Cervical Cancer Screening. Available online: <https://cancerregistry.fi/research/research-themes/cervical-cancer-screening/> (accessed on 5 June 2019).

53. Finnish Institute for Health & Welfare. Care Register for Health Care. Available online: <https://thl.fi/en/web/thlfi-en/statistics/information-on-statistics/register-descriptions/care-register-for-health-care> (accessed on 5 June 2019).
54. Finnish Institute for Health & Welfare. Medical Birth Register. Available online: <https://thl.fi/en/web/thlfi-en/statistics/information-on-statistics/register-descriptions/newborns> (accessed on 5 June 2019).
55. Finnish Institute for Health & Welfare. Description of Statistics-New-Born. Available online: <https://thl.fi/en/web/thlfi-en/statistics/information-on-statistics/description-of-statistics/newborns> (accessed on 5 June 2019).
56. Statistics Finland. Statistics. Available online: https://www.stat.fi/til/index_en.html (accessed on 5 June 2019).
57. The Social Insurance Institution of Finland (Kela). Statistics by Topic. Available online: <https://www.kela.fi/web/en/statistics-by-topic> (accessed on 5 June 2019).
58. Population Finland. Personal Identity Code. Available online: <https://vrk.fi/en/personal-identity-code1> (accessed on 5 June 2019).
59. *Research Triangle Institute: SUDAAN Language Manual, Volumes 1 and 2, Release 11*; Research Triangle Institute: Research Triangle Park, NC, USA, 2012.
60. Akaike, H. Information theory as an extension of the maximum likelihood principle. In *Second International Symposium on Information Theory*; Csáki, F., Petrov, B.N., Eds.; BNPBF Csaki Budapest Akademiai Kiado: Budapest, Hungary, 1973; pp. 267–281.
61. Graubard, B.; Korn, E. Predictive margins with survey data. *Biometrics* **1999**, *55*, 652–659. [[CrossRef](#)] [[PubMed](#)]
62. Lehtonen, R.; Pahkinen, E. *Practical Methods for Design and Analyses of Complex Surveys*, 2nd ed.; John Wiley and Sons: Chichester, UK, 2004.
63. Regulation GDP. Regulation (EU) 2016/679 of the European Parliament and of the Council of 27 April 2016 on the protection of natural persons with regard to the processing of personal data and on the free movement of such data and repealing Directive 95/46. *J. Eur. Union* **2016**, *59*, 294.
64. Moen, K.A.; Kumar, B.; Qureshi, S.; Diaz, E. Differences in cervical cancer screening between immigrants and nonimmigrants in Norway: A primary healthcare register-based study. *Eur. J. Cancer Prev.* **2017**, *26*, 521–527. [[CrossRef](#)] [[PubMed](#)]
65. Finnish Ministry of Social Affairs and Health. Health Services. Available online: <http://stm.fi/en/health-services> (accessed on 9 September 2017).
66. Team, V.; Manderson, L.H.; Markovic, M. From state care to self-care: Cancer screening behaviours among Russian-speaking Australian women. *Aust. J. Prim. Health* **2013**, *19*, 130–137. [[CrossRef](#)] [[PubMed](#)]
67. Carballo, M.; Nerukar, A. Migration, refugees, and health risks. *Emerg Infect Dis.* **2001**, *7* (Suppl. S3), 556–600. [[CrossRef](#)] [[PubMed](#)]
68. De Sanjosé, S.; Serrano, B.; Castellsagué, X.; Brotons, M.; Muñoz, J.; Bruni, L.; Bosch, F.X. Human papillomavirus (HPV) and related cancers in the global Alliance for vaccines and immunization (GAVI) countries: A WHO/ICO HPV information Centre report. *Vaccine* **2012**, *30* (Suppl. S4). [[CrossRef](#)]
69. Anttila, A.; Nieminen, P. Cervical cancer screening programme in Finland with an example on implementing alternative screening methods. *Coll. Antropol.* **2007**, *31*, 17–22.
70. Chang, H.K.; Seo, S.; Myong, J.; Koo, J.; Jeong, J. Factors Associated with Cervical Cancer Screening among Married Female Immigrants with Korean Husbands in South Korea. *Int. J. Environ. Res. Public Health* **2018**, *15*, 2528. [[CrossRef](#)] [[PubMed](#)]
71. Castaneda, A.; Rask, S.; Koponen, P.; Mölsä, M.; Koskinen, S.; Migrant Health and Wellbeing. A Study on Persons of Russian, Somali and Kurdish Origin in Finland. *Report 61. (In Finnish, English abstract). Helsinki, Finland. Finnish Institute for Health & Welfare (THL)*. Available online: <http://urn.fi/URN:ISBN:978-952-245-739-4> (accessed on 1 May 2018).
72. Willems, B.; Bracke, P. The education gradient in cancer screening participation: A consistent phenomenon across Europe? *Int. J. Public Health* **2018**, *63*, 93–103. [[CrossRef](#)]
73. Lorini, C.; Caini, S.; Ierardi, F.; Bachini, L.; Gemmi, F.; Bonaccorsi, G. Health Literacy as a Shared Capacity: Does the Health Literacy of a Country Influence the Health Disparities among Immigrants? *Int. J. Environ. Res. Public Health* **2020**, *17*, 1149. [[CrossRef](#)] [[PubMed](#)]
74. The Social Insurance Institution of Finland (Kela). Old-Age Pension. Available online: <https://www.kela.fi/web/en/old-age-pension?inheritRedirect=true> (accessed on 8 October 2019).

75. Sund, R. Quality of the Finnish Hospital Discharge Register: A systematic review. *Scand. J. Public Health* **2012**, *40*, 505–515. [[CrossRef](#)] [[PubMed](#)]
76. Gissler, M.; Haukka, J. Finnish health and social welfare registers in epidemiological research. *Nor. Epidemiol.* **2004**, *14*, 113–120. [[CrossRef](#)]
77. Pukkala, E.; Engholm, G.; Højsgaard Schmidt, L.K.; Storm, H.; Khan, S.; Lambe, M.; Pettersson, D.; Ólafsdóttir, E.; Tryggvadóttir, L.; Hakanen, T.; et al. Nordic Cancer Registries—An overview of their procedures and data comparability. *Acta Oncol.* **2018**, *57*, 440–455. [[CrossRef](#)]
78. Lofters, A.; Vahabi, M.; Glazier, R.H. The validity of self-reported cancer screening history and the role of social disadvantage in Ontario, Canada. *BMC Public Health* **2015**, *15*, 28. [[CrossRef](#)]

Publisher’s Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



© 2020 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).