

2018

# Body fat mass, lean body mass and associated biomarkers as determinants of bone mineral density in children 6-8 years of age - The Physical Activity and Nutrition in Children (PANIC) study

Soininen, Sonja

Elsevier BV

---

article

info:eu-repo/semantics/acceptedVersion

© Elsevier Inc.

CC BY-NC-ND <https://creativecommons.org/licenses/by-nc-nd/4.0/>

<http://dx.doi.org/10.1016/j.bone.2018.01.003>

---

<https://erepo.uef.fi/handle/123456789/6125>

*Downloaded from University of Eastern Finland's eRepository*

# Body fat mass, lean body mass and associated biomarkers as determinants of bone mineral density in children 6–8 years of age – The Physical Activity and Nutrition in Children (PANIC) Study

Sonja Soininen<sup>a,b,c,\*</sup>, Virpi Sidoroff<sup>d</sup>, Virpi Lindi<sup>a</sup>, Anitta Mahonen<sup>e</sup>, Liisa Kröger<sup>f</sup>, Heikki Kröger<sup>g,h</sup>, Jarmo Jääskeläinen<sup>f</sup>, Mustafa Atalay<sup>a</sup>, David E. Laaksonen<sup>a,i</sup>, Tomi Laitinen<sup>j</sup>, Timo A. Lakka<sup>a,j,k</sup>

Affiliations:

<sup>a</sup> Institute of Biomedicine, Physiology, School of Medicine, University of Eastern Finland, PO Box 1627, 70211 Kuopio, Finland

<sup>b</sup> Institute of Dentistry, University of Eastern Finland, PO Box 1627, 70211 Kuopio, Finland

<sup>c</sup> Social and Health Center, City of Varkaus, Savontie 55, 78300 Varkaus, Finland

<sup>d</sup> Department of Pediatrics, North-Karelia Central Hospital, Tikkamäentie 16, 80210 Joensuu, Finland

<sup>e</sup> Institute of Biomedicine, Medical Biochemistry, School of Medicine, University of Eastern Finland, PO Box 1627, Kuopio, Finland

<sup>f</sup> Department of Pediatrics, Kuopio University Hospital and University of Eastern Finland, PO Box 100, 70029 Kuopio, Finland

<sup>g</sup> Department of Orthopedics and Traumatology, Kuopio University Hospital, PO Box 100, 70029 Kuopio, Finland.

<sup>h</sup> Kuopio Musculoskeletal Research Unit (KMRU), University of Eastern Finland, PO Box 1627, 70211 Kuopio, Finland.

<sup>i</sup> Department of Internal Medicine, Kuopio University Hospital, PO Box 100, 70029 Kuopio, Finland

<sup>j</sup> Department of Clinical Physiology and Nuclear Medicine, Kuopio University Hospital, PO Box 100, 70029 Kuopio, Finland

<sup>k</sup> Kuopio Research Institute of Exercise Medicine, Haapaniementie 16, 70100 Kuopio, Finland

**\*Corresponding author:** Sonja Soininen, University of Eastern Finland, Institute of Biomedicine / Physiology, PO Box 1627, Fin-70211 Kuopio, Finland; [sonja.soininen@uef.fi](mailto:sonja.soininen@uef.fi)

**e-mail addresses:** [sonja.soininen@uef.fi](mailto:sonja.soininen@uef.fi); [virpi.sidoroff@siunsoite.fi](mailto:virpi.sidoroff@siunsoite.fi); [virpi.lindi@uef.fi](mailto:virpi.lindi@uef.fi); [anitta.mahonen@uef.fi](mailto:anitta.mahonen@uef.fi); [liisa.kroger@kuh.fi](mailto:liisa.kroger@kuh.fi); [heikki.kroger@kuh.fi](mailto:heikki.kroger@kuh.fi); [jarmo.jaaskelainen@uef.fi](mailto:jarmo.jaaskelainen@uef.fi); [mustafa.atalay@uef.fi](mailto:mustafa.atalay@uef.fi); [david.laaksonen@uef.fi](mailto:david.laaksonen@uef.fi); [tomi.laitinen@kuh.fi](mailto:tomi.laitinen@kuh.fi); [timo.lakka@uef.fi](mailto:timo.lakka@uef.fi)

37    **Abstract**

38    Lean body mass (LM) has been positively associated with bone mineral density (BMD) in children  
39    and adolescents, but the relationship between body fat mass (FM) and BMD remains controversial.  
40    Several biomarkers secreted by adipose tissue, skeletal muscle, or bone may affect bone metabolism  
41    and BMD. We investigated the associations of LM, FM, and such biomarkers with BMD in children.

42  
43    We studied a population sample of 472 prepubertal Finnish children (227 girls, 245 boys) aged 6-8  
44    years. We assessed BMD, LM, and FM using whole-body dual-energy x-ray absorptiometry and  
45    analysed several biomarkers from fasting blood samples. We studied the associations of LM, FM,  
46    and the biomarkers with BMD of the whole body excluding the head using linear regression analysis.

47  
48    LM (standardized regression coefficient  $\beta=0.708$ ,  $p<0.001$ ), FM ( $\beta=0.358$ ,  $p<0.001$ ), and irisin  
49    ( $\beta=0.079$ ,  $p=0.048$ ) were positive correlates for BMD adjusted for age, sex, and height in all children.  
50    These associations remained statistically significant after further adjustment for LM or FM. The  
51    positive associations of dehydroepiandrosterone sulphate (DHEAS), insulin, homeostatic model  
52    assessment for insulin resistance (HOMA-IR), leptin, free leptin index, and high-sensitivity C-  
53    reactive protein and the negative association of leptin receptor with BMD were explained by FM. The  
54    positive associations of DHEAS and HOMA-IR with BMD were also explained by LM. Serum 25-  
55    hydroxyvitamin D was a positive correlate for BMD adjusted for age, sex, and height and after further  
56    adjustment for FM but not for LM. LM and FM were positive correlates for BMD also in girls and  
57    boys separately. In girls, insulin, HOMA-IR, leptin, and free leptin index were positively and leptin  
58    receptor was negatively associated with BMD adjusted for age, height, and LM. After adjustment for  
59    age, height, and FM, none of the biomarkers was associated with BMD. In boys, leptin and free leptin  
60    index were positively and leptin receptor was negatively associated with BMD adjusted for age,  
61    height, and LM. After adjustment for age, height and FM, 25(OH)D was positively and IGF-1 and  
62    leptin were negatively associated with BMD. FM strongly modified the association between leptin  
63    and BMD.

64  
65    LM but also FM were strong, independent positive correlates for BMD in all children, girls, and boys.  
66    Irisin was positively and independently associated with BMD in all children. The associations of other  
67    biomarkers with BMD were explained by LM or FM.

68    **Keywords:** bone mineral density; lean body mass; body fat mass; DXA; child; cytokine

69    **Abbreviations**

70    BF%, body fat percentage

71    BMC, bone mineral content

72    BMD, bone mineral density

73    BMI, body mass index

74    DHEAS, dehydroepiandrosterone sulphate

75    DXA, dual-energy x-ray absorptiometry

76    FM, body fat mass

77    HOMA-IR, the homeostatic model assessment for insulin resistance

78    hs-CRP, high-sensitivity C-reactive protein

79    IGF-1, insulin-like growth factor 1

80    IL-6, interleukin 6

81    LM, lean body mass

82    SD, standard deviation

83    SDS, standard deviation score

84    TNF- $\alpha$ , tumor necrosis factor  $\alpha$

85    25(OH)D, 25-hydroxyvitamin D

## 1. Introduction

Early childhood and puberty are the periods of rapid growth and bone accretion, and the majority of bone mass is gained during adolescence and early adulthood [1–3]. Bone mineral accrual during growth is dependent on multiple factors such as genetic background, sex, race, nutrition, physical activity, and hormone metabolism [2,3]. Higher lean body mass (LM) has been associated with higher bone mineral density (BMD) and bone mineral content (BMC) in children and adolescents [4–7], but the relationship of body fat mass (FM) with BMD or BMC remains controversial [5,6,8–10]. FM has been positively associated with BMD independent of LM in prepubertal children [6]. However, there is some evidence that higher FM is detrimental to bone accrual during and after puberty [5,8,9] and that overweight children and adolescents are at an increased risk of forearm fractures [10].

Mechanical loading increases bone formation, and weight-bearing exercise improves bone mineral accrual [11]. The classical Wolff's law and later the Frost's mechanostat theory propose that bone strength is regulated by modeling and remodeling processes which depend on the forces acting on the bones [12]. The mechanical load to bone is increased not only because of physical activity and increased muscle mass but also due to increased FM and particularly obesity [3].

In addition to the mechanical load, adipose tissue may influence bone metabolism through adipokines, other cytokines, and hormones [13–15]. Adipose tissue may stimulate bone formation by producing estrogens from steroid precursors and by increasing circulating leptin and insulin levels [13–15]. However, adipose tissue produces adiponectin and inflammation-related cytokines, such as tumor necrosis factor  $\alpha$  (TNF- $\alpha$ ) and interleukin 6 (IL-6), which may have deleterious effects on bone [13–15]. Vitamin D is a prohormone converted in the liver to 25-hydroxyvitamin D (25[OH]D) and then in the kidney to 1,25-dihydroxyvitamin D (1,25[OH]<sup>2</sup>D), the active metabolite which regulates calcium, phosphorus, and bone metabolism [16]. Obesity has been associated with lower serum levels of 25(OH)D [17], that could therefore be one of the links between obesity and BMD.

More recently, also skeletal muscle and bone have been recognized as endocrine organs [18,19]. Skeletal muscle produces myokines, such as myostatin, insulin-like growth factor I (IGF-1), irisin, and IL-6, which may be important mediators in the interaction between skeletal muscle and bone [18,19]. IGF-1 may be one of the factors that mediate the response of bone and skeletal muscle to mechanical loading [19,20]. Osteocytes also secrete IL-6, IGF-1, and other hormone-like factors,

118 such as osteocalcin and fibroblast growth factor 23, which have been suggested to play a role in the  
119 association between skeletal muscle and bone metabolism [18,19].

120

121 Low BMD in childhood tends to persist until young adulthood [21], and bone mass attained during  
122 childhood and adolescence is one of the most important determinants of lifelong skeletal health [22].  
123 Pediatric obesity is a growing global health problem [23], and it is therefore important to know how  
124 adiposity and associated increase in LM affects BMD among children. There is no consensus on the  
125 associations of FM and LM with BMD or the underlying mechanisms. We therefore studied the  
126 associations of LM, FM, and associated biomarkers, including adipokines, myokines, inflammation-  
127 related biomarkers, growth factors, and 25(OH)D, with BMD assessed by dual-energy x-ray  
128 absorptiometry (DXA) in a population sample of children 6-8 years of age.

## 129 **2. Methods**

### 130 *2.1 Study design and participants*

131 The present analyses are based on the baseline data of the Physical Activity and Nutrition in Children  
132 (PANIC) Study, which is an ongoing physical activity and dietary intervention study in a population  
133 sample of children 6–8 years of age from the city of Kuopio, Finland (ClinicalTrials.gov registration  
134 number NCT01803776). Altogether 736 children from the primary schools of Kuopio were invited  
135 to participate in the baseline examinations in 2007–2009. Of the invited children, 512 (70%)  
136 participated in the baseline examinations. The participants did not differ in age, sex distribution, or  
137 body mass index standard deviation score (BMI-SDS) from all children who started the 1<sup>st</sup> grade in  
138 the city of Kuopio in 2007–2009 based on data from the standard school health examinations. From  
139 the present analyses, we excluded children who had chronic diseases or medications that could affect  
140 BMD, such as juvenile arthritis demanding long-term treatment with oral corticosteroids. We also  
141 excluded 12 children who had entered puberty to avoid associated confounding. Complete data on  
142 the main variables used in the present analyses were available for 472 children (227 girls, 245 boys).  
143 The study was conducted according to the ethical guidelines laid down in the Declaration of Helsinki.  
144 The study protocol was approved by the Research Ethics Committee of the Hospital District of  
145 Northern Savo. Both children and their parents gave their written informed consent.

## 146 2.2 Assessment of bone mineral density and body composition

147 LM, FM, body fat percentage (BF %), and BMD of the whole body excluding the head were assessed  
148 using the Lunar Prodigy Advance<sup>®</sup> DXA device (GE Medical Systems, Madison, WI, USA) and the  
149 Encore<sup>®</sup> software, Version 10.51.006 (GE Company, Madison, WI, USA), according to the  
150 manufacturer's instructions using standardized protocols. The same DXA device and software were  
151 used in all measurements. Body weight was measured twice the children having fasted for 12 hours,  
152 emptied the bladder, and standing in light underwear by the InBody<sup>®</sup> 720 bioelectrical impedance  
153 device (Biospace, Seoul, Korea) to accuracy of 0.1 kg. The mean of these two values was used in the  
154 analyses. Body height was measured three times the children standing in the Frankfurt plane without  
155 shoes using a wall-mounted stadiometer to accuracy of 0.1 cm. The mean of the nearest two values  
156 was used in the analyses. BMI-SDS was calculated using national reference values [24]. Waist  
157 circumference was measured three times after expiration at mid-distance between the bottom of the  
158 rib cage and the top of the iliac crest with an unstretchable measuring tape to accuracy of 0.1 cm. The  
159 mean of the nearest two values was used in the analyses. Intraclass correlation coefficients for body  
160 weight and height and waist circumference were >0.99.

## 161 2.3 Biochemical analyses

162 Venous blood samples were taken the children having fasted for 12 hours. Blood was immediately  
163 centrifuged and stored at a temperature of -75°C until biochemical analyses, except for glucose that  
164 was measured from non-frozen plasma samples. Serum 25(OH)D concentration was analysed by a  
165 chemiluminescence immunoassay called the LIAISON<sup>®</sup> 25 OH Vitamin D TOTAL Assay (DiaSorin  
166 Inc., Stillwater, USA) as described earlier [25,26]. Serum dehydroepiandrosterone sulphate (DHEAS)  
167 concentration was used as a marker of biochemical adrenarche and was determined using an enzyme  
168 linked immunosorbent assay (ELISA) kit (Alpha Diagnostic International, San Antonio, Texas, USA)  
169 [27]. Serum IGF-1 concentration was analysed using an ELISA kit (Mediagnost, Reutlingen,  
170 Germany). Plasma glucose concentration was measured using the hexokinase method (Roche  
171 Diagnostics GmbH, Mannheim, Germany). Serum insulin concentration was measured by the  
172 electrochemiluminescence immunoassay with the sandwich principle (Roche Diagnostics GmbH,  
173 Mannheim, Germany). We calculated the Homeostatic Model Assessment for Insulin Resistance  
174 (HOMA-IR) using the formula fasting serum insulin x fasting plasma glucose/22. Serum high-  
175 molecular-weight adiponectin concentration was analysed using an ELISA kit after a specific  
176 proteolytic digestion of other multimeric adiponectin forms (Millipore, Billerica, MA, USA). Plasma  
177 leptin concentration was measured by a competitive radioimmunoassay (Multigamma 1261-001,

178 PerkinElmer Wallac Oy, Turku, Finland) and plasma soluble leptin receptor concentration using an  
179 ELISA kit (Multicalc evaluation programme PerkinElmer Wallac Oy, Turku, Finland). We calculated  
180 the free leptin index by dividing leptin with soluble leptin receptor and multiplying by 100 [28].  
181 Commercially available ELISA kits were employed for the measurement of plasma irisin (Phoenix  
182 Pharmaceuticals, Burlingame, California, USA), IL-6, and TNF- $\alpha$  concentrations (Sanquin Reagents,  
183 Amsterdam, The Netherlands). Plasma high-sensitivity C-reactive protein (hsCRP) was measured  
184 using an enhanced immunoturbidimetric assay with the CRP (Latex) High Sensitive Assay reagent  
185 (Roche Diagnostics GmbH, Mannheim, Germany) and the limit of quantitation of 0.3 mg/l.

#### 186 *2.4 Assessments of general health, puberty, and adrenarche*

187 The parents filled out a questionnaire that included items on the children's chronic diseases and  
188 allergies diagnosed by a physician as well as detailed information on the children's use of  
189 medications. A research physician assessed pubertal status during a medical examination. Central  
190 puberty was defined as breast development at Tanner stage  $\geq 2$  for girls and testicular volume  $\geq 4$  mL  
191 assessed using an orchidometer for boys. Premature adrenarche was defined as serum DHEAS  $\geq 1$   
192  $\mu\text{mol/l}$  ( $\geq 37 \mu\text{g/dl}$ ) [29] and at least one clinical sign of androgen action. Birth weight was obtained  
193 from Kuopio University Hospital record, and birth weight -SDS was calculated according to Finnish  
194 growth reference data [30].

#### 195 *2.5 Statistical methods*

196 We performed statistical analyses using the IBM SPSS Statistics<sup>®</sup> software, Version 21 (IBM Corp.,  
197 Armonk, NY, USA). The normality of distributions of the variables was verified visually and by the  
198 Kolmogorov-Smirnov test. The t-test for independent samples and the Mann-Whitney's U-test were  
199 used to examine differences in the basic characteristics between sexes. Linear regression analysis was  
200 used to investigate the determinants of BMD, and the normality of residuals for regression models  
201 was assessed using histograms. Model 1 included each determinant of BMD separately, adjusted for  
202 age and sex. Model 2 was additionally adjusted for body height. Model 3 included all variables in  
203 Model 2 and LM, and Model 4 included all variables in Model 2 and FM. Corresponding linear  
204 regression analyses were also performed for girls and boys separately. FM had a strong positive  
205 correlation with leptin in girls ( $r=0.789$ ,  $p<0.001$ ), boys ( $r=0.850$ ,  $p<0.001$ ), and girls and boys  
206 combined ( $r=0.810$ ,  $p<0.001$ ). We therefore tested whether FM modified the association between  
207 leptin and BMD by analyzing this association in the sex-specific thirds of FM using linear regression



analysis adjusted for age, sex, and body height. In all analyses, associations with a p-value of <0.05 were considered statistically significant.

**3. Results**

*3.1 Characteristics of children*

The boys were heavier and taller and had higher waist circumference and LM and lower BF% and FM than the girls, but there was no difference in BMI-SDS between the genders (Table 1). The girls had higher IGF-1, insulin, leptin, and free leptin index and lower leptin receptor and IL-6 than the boys. Of the children, 38 (8.1%) had asthma, 128 (27.1%) any allergic symptom (rhinitis, conjunctivitis, atopy, food or medicine allergy), 21 (4.4%) an attention deficit hyperactivity disorder (ADHD/ADD) or another mild neurocognitive disorder or developmental delay, 8 (1.7%) a mild congenital dysmorphism, and 10 (2.1%) any other chronic disease. There was no difference in BMD between children with these diseases and those without them.

*3.2. Determinants of bone mineral density in all children*

Body height ( $\beta=0.572$ ,  $p<0.001$ ) and weight ( $\beta=0.709$ ,  $p<0.001$ ) were positively associated with BMD adjusted for age and sex. LM was also a strong positive correlate for BMD adjusted for age and sex (Table 2, Model 1). This association remained similar after additional adjustment for body height (Model 2) but weakened slightly after further adjustment for FM (Model 4). Moreover, FM had a strong positive association with BMD adjusted for age and sex (Table 2, Model 1). This association weakened after additional adjustment for body height (Model 2) but remained similar when further adjusted for LM (Model 3). Birth weight was positively associated with BMD adjusted for age and sex (Table 2, Model 1), but this association disappeared after additional adjustments (Models 2-4).

Serum 25(OH)D was positively associated with BMD adjusted for age and sex (Table 2, Model 1). This association remained almost similar after additional adjustment for body height and FM (Models 2 and 4) but was no longer statistically significant when adjusted for LM (Model 3). DHEAS was positively associated with BMD adjusted for age and sex (Table 2, Model 1). This association weakened when additionally adjusted for body height (Model 2) but was no longer statistically significant after adjustment for LM or FM (Models 3-4). IGF-1 was a positive correlate for BMD adjusted for age and sex (Table 2, Model 1) but not after further adjustments (Models 2-4). Insulin and HOMA-IR were positively associated with BMD adjusted for age and sex (Table 2, Model 1).

These associations weakened after additional adjustment for body height (Model 2). The association of insulin weakened and that of HOMA-IR was no longer statistically significant after further adjustment for LM (Model 3). The associations of insulin and HOMA-IR with BMD disappeared when adjusted for FM (Model 4).

Adiponectin was a negative correlate for BMD adjusted for age and sex (Table 2, Model 1) but not after further adjustments (Models 2-4). Leptin was positively associated with BMD adjusted for age and sex (Table 2, Model 1). This association weakened after additional adjustment for body height and LM (Models 2-3) and was no longer statistically significant after adjustment for FM (Model 4). There was a positive association between leptin and BMD in the highest sex-specific third of FM ( $\beta=0.274$ ,  $p<0.001$ ) but a non-significant inverse association in the middle third ( $\beta=-0.144$ ,  $p=0.058$ ) and the lowest third ( $\beta=-0.112$ ,  $p=0.118$ ) adjusted for age and body height. Lower leptin receptor and higher free leptin index were associated with higher BMD adjusted for age and sex (Table 2, Model 1). These associations weakened after additional adjustment for body height and when further adjusted for LM (Models 2-3) and were no longer statistically significant after adjustment for FM (Model 4). Irisin was positively associated with BMD adjusted for age and sex (Table 2, Model 1). This association weakened slightly when additionally adjusted for body height (Model 2) and remained similar after further adjustment for LM or FM (Models 3-4).

IL-6 and TNF- $\alpha$  were not associated with BMD (Table 2, Models 1-4). Higher hs-CRP was associated with higher BMD adjusted for age and sex (Table 2, Model 1), after additional adjustment for body height (Model 2), and also when further adjusted for LM (Model 3). However, this association disappeared after adjustment for FM (Model 4).

### *3.2.2 Determinants of bone mineral density in girls*

In girls, body height ( $\beta=0.615$ ,  $p<0.001$ ) and weight ( $\beta=0.727$ ,  $p<0.001$ ) were positively associated with BMD adjusted for age. LM had a strong positive association with BMD adjusted for age, body height, and FM (Table 3, Models 1, 2, and 4). FM was also a strong positive correlate for BMD adjusted for age, body height, and LM (Table 3, Models 1-3). Birth weight SDS, 25(OH)D, DHEAS, IGF-1, and irisin were positively associated with BMD when adjusted for age (Table 3, Model 1) but not after further adjustments (Models 2-4). Insulin and HOMA-IR were positive correlates for BMD adjusted for age, body height, and LM (Table 3, Models 1-3) but not when adjusted for FM (Model 4). Leptin and free leptin index were positively and leptin receptor was negatively associated with BMD adjusted for age, body height, and LM (Table 3, Models 1-3) but not adjusted for FM (Model

271 4). There was a positive association between leptin and BMD in the highest third of FM ( $\beta=0.346$ ,  
272  $p<0.001$ ) but a non-significant inverse association in the middle third ( $\beta=-0.169$ ,  $p=0.126$ ) and the  
273 lowest third ( $\beta=-0.122$ ,  $p=0.261$ ) adjusted for age and body height.

### 274 3.2.3 Determinants of bone mineral density in boys

275 In boys, body height ( $\beta=0.520$ ,  $p<0.001$ ) and weight ( $\beta=0.686$ ,  $p<0.001$ ) were positively associated  
276 with BMD adjusted for age. LM had a strong positive association with BMD adjusted for age, body  
277 height, and FM (Table 4, Models 1, 2, and 4). FM was also a strong positive correlate for BMD  
278 adjusted for age, body height, and LM (Table 4, Models 1-3). Serum 25(OH)D was positively  
279 associated with BMD adjusted for age, body height, and FM (Table 4, Models 2 and 4) but not  
280 adjusted for LM (Model 4). Birth weight SDS, DHEAS, insulin, HOMA-IR and hs-CRP were  
281 positively associated with BMD adjusted for age (Table 4, Model 1) but not after further adjustments  
282 (Models 2-4). IGF-1 was negatively associated with BMD only when adjusted for age, body height,  
283 and FM (Table 4, Model 4). Leptin and free leptin index were positively and leptin receptor was  
284 negatively associated with BMD adjusted for age, body height, and LM (Table 4, Models 1-3), but  
285 the associations of free leptin index and leptin receptor were no longer statistically significant and  
286 that of leptin became negative when adjusted for LM (Model 4). There was a non-significant positive  
287 association between leptin and BMD in the highest third of FM ( $\beta=0.199$ ,  $p=0.061$ ), a non-significant  
288 inverse association in the middle third ( $\beta=-0.135$ ,  $p=0.203$ ) and no association in the lowest third ( $\beta=-$   
289  $0.024$ ,  $p=0.821$ ).

## 290 4. Discussion

291 Our study is one of the few studies on the associations of LM, FM, and various biomarkers secreted  
292 by adipose tissue, skeletal muscle, or bone with BMD in a population sample of prepubertal children.  
293 LM but also FM were strong and independent positive determinants of BMD in all children, girls,  
294 and boys. Plasma irisin was also an independent positive correlate for BMD in all children but not in  
295 girls and boys separately. The associations of other biomarkers were explained by body height, LM,  
296 or FM. In boys, the positive association between leptin and BMD became negative and the negative  
297 association between IGF-1 and BMD strengthened after controlling for FM.

298

299 In line with previous studies among children and adolescents [4,5,7], LM was a strong positive  
300 correlate for BMD in the current study. The positive association between LM and BMD may be

301 explained by increased mechanical load to bone caused by increased LM and the loading effect of  
302 weight-bearing exercise on bone mass and metabolism [11].  
303

304 A recently identified myokine irisin is produced by skeletal muscle after exercise and may increase  
305 energy expenditure [31]. Irisin has been found to increase bone mass in mice [32], but evidence on  
306 the association between serum irisin and BMD in humans is limited. Irisin has been positively  
307 associated with bone mass and strength in young athletes and negatively related to vertebral fragility  
308 fractures in postmenopausal women [31,33]. To the best of our knowledge, the association between  
309 irisin and BMD has not been studied earlier in children. We found that higher serum irisin levels were  
310 associated with higher BMD even after controlling for LM or FM. The weak positive association  
311 between irisin and BMD was slightly stronger in girls than in boys, but statistical power was limited  
312 in these sex-specific analyses.  
313

314 Of other biomarkers previously related to skeletal muscle and bone metabolism, insulin had a weak  
315 positive association with BMD even after controlling for LM. However, the association between  
316 insulin and BMD was explained by FM. IGF-1 was positively associated with BMD in all children  
317 and in girls but not after controlling for body size and composition. Moreover, there was a weak  
318 negative association between IGF-1 and BMD in boys when controlled for FM. Previous studies in  
319 children and adolescents have reported an independent positive association between IGF-1 and bone  
320 growth [20] and a muscle-dependent positive association between IGF-1 and BMD [20,34]. However,  
321 insulin resistance has suppressed the muscle-dependent relationship between IGF-1 and BMC and  
322 cortical bone measurements in children 9-13 years of age [34,35]. One reason for the inconsistency  
323 between our results and the findings of earlier studies could be that our participants were prepubertal  
324 and slightly younger than those of the previous studies. It is also possible that the weak negative  
325 association between IGF-1 and BMD in boys after controlling for FM in our study is partly explained  
326 by the positive relationships among adiposity, insulin resistance, and IGF-1.  
327

328 FM has been positively associated with BMD in some previous studies among mainly prepubertal  
329 children [6,36]. Obesity has also been associated with increased bone mass independent of LM in a  
330 study among children and adolescents [37]. Moreover, adiposity was associated with increased bone  
331 mass in another study in adolescents, but this association was explained by LM [7]. One explanation  
332 for the positive association between FM and BMD among children and adolescents could be the  
333 increased mechanical load to the bone due to adiposity [3]. Another reason could be that adipose  
334 tissue stimulates bone growth [36]. However, one study reported a decreased volumetric BMD in

335 obese prepubertal children despite increased bone size [38]. Another study showed an inverse  
336 association between BF% and BMD in adolescents [5]. In a Finnish study among prepubertal and  
337 pubertal children, those with decreased body fat content and those with increased fat content had  
338 decreased BMD independent of LM [39]. In the current study, FM was positively associated with  
339 BMD independent of LM, even though LM was a stronger correlate for BMD than FM. This  
340 observation is consistent with the results of a previous study among children [6]. Studies that have  
341 shown an association between excess fat mass and decreased BMD have been conducted in older and  
342 more overweight children and adolescents [5,39] than the participants of our study. Only 14% of the  
343 girls and 10% of the boys in our population sample of prepubertal children 6-8 years of age were  
344 overweight or obese [40]. Therefore, we cannot draw a conclusion on the association between obesity  
345 and BMD based on our findings. It is possible that the detrimental effect of excess fat mass appears  
346 in later childhood or in adolescence during or after puberty along with changes in body composition  
347 [1]. In our study, the association between LM and BMD was stronger in boys than in girls. One reason  
348 for this finding could be that boys have more skeletal muscle and girls have more adipose tissue  
349 already in prepubertal stage [1], that is consistent with our observation.

350  
351 Leptin is an adipocyte-secreted hormone that decreases appetite and increases energy expenditure  
352 [14] but may also influence bone modeling through central and peripheral mechanisms [14,15].  
353 Leptin has been suggested to inhibit bone formation indirectly through the sympathetic nervous  
354 system [14,15]. In contrast, leptin directly enhances bone formation and inhibits bone resorption  
355 peripherally, even though the mechanisms are rather complex and not yet well defined [14,15]. These  
356 local effects of leptin on bone have been suggested to be dominant, and higher circulating leptin levels  
357 may therefore be related to a stronger skeleton [15]. Leptin may also regulate the hypothalamic-  
358 pituitary-peripheral endocrine axes, including thyroid, gonadal, cortisol, and growth hormone axes,  
359 which are possible additional indirect ways by which leptin affects bone [41]. Soluble leptin receptor  
360 is the major protein binding leptin in blood, and leptin receptor levels seem to vary independent of  
361 serum leptin levels during childhood [28]. Functional differences between free and bound leptin are  
362 not clear, but some studies have suggested that free leptin index better reflects the physiological  
363 actions of leptin [28]. A meta-analysis concluded that circulating leptin levels were positively  
364 associated with BMD [42], but most of the 46 studies included in the analysis were performed in  
365 adults. The association between leptin and total body BMD was also positive in five studies among  
366 girls [42]. Interestingly, the relationship between leptin and BMD adjusted for body mass was  
367 negative in the only small study among boys [43]. Furthermore, body fat content was not taken into  
368 account in the meta-analysis [42]. In a previous study, free leptin index was associated with bone

turnover markers [13], which may be one mechanism for the inverse association between leptin and BMD. We found that leptin receptor level was negatively and leptin and free leptin index were positively associated with BMD independent of LM, but these associations were explained by FM. Moreover, the association between leptin and BMD became negative in boys after controlling for FM. Leptin was positively associated with BMD in the highest sex-specific third of FM but had a weak negative association in the middle and lowest thirds. These findings suggest that FM strongly modifies the association between leptin and BMD.

Adiponectin is an adipokine that has been inversely related to FM in children [44], and this inverse association has been found to strengthen in puberty [45]. Adiponectin regulates energy homeostasis, glucose and lipid metabolism, and inflammatory pathways [15]. Increased adiponectin has been associated with reduced bone mass in children [44]. This may be explained by the decreased circulating levels of insulin and IGF-1 due to increased adiponectin levels [15]. In the current study among prepubertal children, we found a weak negative association between adiponectin and BMD, but it was largely explained by LM and FM. It is possible that the negative association between adiponectin and BMD might be stronger after puberty.

Excess adiposity is associated with insulin resistance and hyperinsulinemia in youth [46]. Insulin has been suggested to be anabolic for bone formation, and higher serum insulin levels have been associated with higher BMD in adults [15]. However, the associations of insulin resistance with BMC and BMD remain controversial in children and adolescents [47–49]. In a study among prepubertal overweight children, BMC was lower in children with prediabetes than in children without it [47]. In overweight adolescents, increased HOMA-IR was associated with decreased BMD [48]. In another study among adolescents, insulin was positively associated with BMD, but the association was inverse after controlling for FM [49]. In line with these results, we found that higher fasting insulin and HOMA-IR were associated with higher BMD, but the associations became weak negative in boys and disappeared in girls after controlling for FM. These findings suggest that the association between insulin resistance and BMD is largely dependent on adiposity that should be taken into account when interpreting the results.

IL-6 has a double-edged role in bone metabolism as it may stimulate both osteocyte differentiation and osteoclastic bone resorption [19]. IL-6 but also TNF- $\alpha$  are inflammation-related cytokines secreted by adipose tissue, and they may enhance bone resorption [14]. We found no association between IL-6 or TNF- $\alpha$  and BMD in children. One explanation for this may be that the prevalence of

overweight was low in our general population of children, and thus the inflammatory-related effects of these cytokines may have been modest. Higher hs-CRP has been associated with lower BMD in adolescent girls [50] and in overweight children with prediabetes but not in overweight children without it [47]. Inconsistent with these findings, we observed a weak positive association between hs-CRP and BMD in children. The reason for this inconsistency probably is the low proportion of overweight and obese children in our population sample [40]. Moreover, the observed positive association between hs-CRP and BMD was explained by FM. This is an expected result as adiposity is known to be related to systemic low-grade inflammation [51].

The definition of vitamin D deficiency based on serum 25(OH)D concentration varies between 25 and 50 nmol/l and the lower limit for optimal serum 25(OH)D concentration has been suggested to be as high as 75 nmol/l [3,16,52–57]. No consensus exists on the optimal serum level of 25(OH)D. As vitamin D is essential for bone metabolism [16], the positive association of 25(OH)D with BMD in the current study was expected, and this is in line with the results of previous studies [4]. However, the association between 25(OH)D and BMD was weak especially in girls, but this is probably explained by the low proportion of children having 25(OH)D concentrations below 50 nmol/l [25], which has been considered as a limit of deficiency based on bone outcomes [53]. The association between 25(OH)D and BMD was stronger in boys, and it was partly explained by LM. One explanation for this finding may be that physically active children, particularly boys, have increased LM and spend more time outdoors and are therefore exposed to sunlight that increases serum 25(OH)D concentrations.

DHEAS is an androgen precursor produced mainly by the adrenal cortex and whose circulating levels are increased during adrenarche [27]. Both obesity and premature adrenarche are associated with advanced bone age [58,59]. However, there are little and inconsistent data on the association between DHEAS and BMD in children [58,60]. In the current study among prepubertal children, higher DHEAS was associated with higher BMD. However, the positive association weakened after controlling for body height, LM, and FM, suggesting that DHEAS does not have an independent effect on BMD in prepubertal children.

Some diseases, conditions and medications, such as juvenile arthritis, renal insufficiency, inflammatory conditions, disabilities, immobility, oral corticosteroid use, or certain antiepileptic drugs, may decrease BMD [61]. We therefore excluded children who had such diseases, conditions, or medications to avoid associated confounding. The use of inhaled corticosteroids has been

437 associated with decreased BMD in some studies [62]. However, a recent review and meta-analysis  
438 concluded that the use of inhaled corticosteroids was not associated with decreased lumbar BMD or  
439 increased risk of fractures [63]. In our study, about 8% of the children had asthma, a few of them used  
440 regular inhaled corticosteroids, and they had similar BMD to children without asthma. We therefore  
441 included children with asthma in the current study population.

442

443 Body weight and BMI have been directly associated with BMD in children and adolescents [3,6], but  
444 neither of them is a specific measure of LM or FM. We therefore investigated the associations of LM  
445 and FM measured by DXA with BMD among children. DXA is also the most widely used method to  
446 evaluate BMD and it has been reported to be well reproducible also in children [64–66]. The  
447 assessment and interpretation of BMD measurements are not simple in growing children because of  
448 both methodological aspects and differences in maturation and growth. In children, The International  
449 Society of Clinical Densitometry (ISCD) recommends measuring BMD and BMC from total body  
450 excluding the head and the posterior-anterior spine [66]. Areal BMD measurements may  
451 underestimate the BMD of short children and overestimate the BMD of tall children. Therefore, ISCD  
452 recommends adjusting BMD of total body excluding head and spinal BMD using height z-score. We  
453 used DXA of the whole body, excluding the head, which is one of the methods recommended to be  
454 used for measuring BMD among children by the ISCD [66]. Moreover, we adjusted the data first for  
455 age and sex and then additionally for body height, all components of height z-score. However, we did  
456 not measure volumetric BMD but areal BMD and did not use computed tomography to measure the  
457 more detailed quality of the bone.

458

459 The results of different studies depend not only on the methods used but also on the age and  
460 maturation of the participants and the prevalence of overweight in the study population, because each  
461 of them affects BMD. We investigated a general population of prepubertal children 6-8 years of age  
462 with a low prevalence of overweight, whereas many other studies have mainly included overweight  
463 or obese children and adolescents with advanced puberty [5,7,37,39,47]. It is therefore difficult to  
464 compare the findings of our study with those of many other studies.

## 465 **5. Conclusions**

466 Our study showed that LM is the strongest positive determinant of BMD, but also FM is positively  
467 and independently associated with BMD in a population sample of mainly normal-weight prepubertal  
468 Finnish children. Of biomarkers related to body composition, irisin had a positive association with  
469 BMD independent of LM and FM. To the best of our knowledge, this is the first study to examine the



association between irisin and BMD in children, and this finding needs to be confirmed in other populations. As expected, 25(OH)D was a positive correlate for BMD, but the association was weak probably due to the relatively low prevalence of vitamin D deficiency in our study population and was partly explained by body composition. In boys, the positive association of leptin with BMD became negative after controlling for FM. This finding suggests that FM strongly modifies the association between leptin and BMD and that adiposity should be taken into account when interpreting the associations of leptin with bone structure and metabolism.

**6. Acknowledgements**

The authors are grateful to all the children and their parents for participating in the PANIC study. The authors are also indebted to the members of the PANIC research team for their skillful contribution in performing the study. The authors are grateful to Ayhan Korkmaz for performing irisin measurements, Leila Antikainen for performing DHEAS and IGF-1 measurements, Tuomas Onnukka for performing leptin measurements, and Kaija Kettunen for performing leptin receptor and adiponectin measurements. We also thank Tarja Kokkola for the help with methodological issues on laboratory measurements.

**7. Funding sources**

This work was financially supported by grants from Ministry of Social Affairs and Health of Finland, Ministry of Education and Culture of Finland, Finnish Innovation Fund Sitra, Social Insurance Institution of Finland, Finnish Cultural Foundation, Juho Vainio Foundation, Foundation for Paediatric Research, Doctoral Programs in Public Health, Paavo Nurmi Foundation, Paulo Foundation, Diabetes Research Foundation, Yrjö Jahnsson Foundation, Finnish Foundation for Cardiovascular Research, Research Committee of the Kuopio University Hospital Catchment Area (State Research Funding), Kuopio University Hospital (previous state research funding (EVO), funding number 5031343), and the city of Kuopio.

**8. Conflict of interest**

The authors declare there are no conflicts of interest.

## 496 9. References

- 497 [1] L.A. Loomba-Albrecht, D.M. Styne, Effect of puberty on body composition, *Curr. Opin.*  
498 *Endocrinol. Diabetes. Obes.* 16 (2009) 10–15. doi:10.1097/MED.0b013e328320d54c.
- 499 [2] A.B. Sopher, I. Fennoy, S.E. Oberfield, An update on childhood bone health: mineral accrual,  
500 assessment and treatment, *Curr. Opin. Endocrinol. Diabetes. Obes.* 22 (2015) 35–40.  
501 doi:10.1097/MED.0000000000000124.
- 502 [3] N.H. Golden, S.A. Abrams, Optimizing bone health in children and adolescents, *Pediatrics.*  
503 134 (2014) e1229–e1243. doi:10.1542/peds.2014-2173.
- 504 [4] M. Pekkinen, H. Viljakainen, E. Saarnio, C. Lamberg-Allardt, O. Mäkitie, Vitamin D is a  
505 major determinant of bone mineral density at school age, *PLoS One.* 7 (2012) e40090.  
506 doi:10.1371/journal.pone.0040090.
- 507 [5] L.N. Mosca, T.B.L. Goldberg, V.N. da Silva, C.C. da Silva, C.S. Kurokawa, A.C. Bisi Rizzo,  
508 J.E. Corrente, Excess body fat negatively affects bone mass in adolescents, *Nutrition.* 30  
509 (2014) 847–852. doi:10.1016/j.nut.2013.12.003.
- 510 [6] M. Heidemann, R. Holst, A.J. Schou, H. Klakk, S. Husby, N. Wedderkopp, C. Molgaard, The  
511 influence of anthropometry and body composition on children’s bone health: the childhood  
512 health, activity and motor performance school (the CHAMPS) study, Denmark, *Calcif. Tissue*  
513 *Int.* 96 (2015) 97–104. doi:10.1007/s00223-014-9941-9.
- 514 [7] L. Gracia-Marco, F.B. Ortega, D. Jiménez-Pavón, G. Rodríguez, M.J. Castillo, G. Vicente-  
515 Rodríguez, L.A. Moreno, Adiposity and bone health in Spanish adolescents. The HELENA  
516 study., *Osteoporos. Int.* 23 (2012) 937–47. doi:10.1007/s00198-011-1649-3.
- 517 [8] P. Dimitri, N. Bishop, J.S. Walsh, R. Eastell, Obesity is a risk factor for fracture in children  
518 but is protective against fracture in adults: A paradox, *Bone.* 50 (2012) 457–466.  
519 doi:10.1016/j.bone.2011.05.011.
- 520 [9] H.A. Weiler, L. Janzen, K. Green, J. Grabowski, M.M. Seshia, K.C. Yuen, Percent body fat  
521 and bone mass in healthy Canadian females 10 to 19 years of age, *Bone.* 27 (2000) 203–207.  
522 doi:10.1016/S8756-3282(00)00314-8.
- 523 [10] A. Goulding, A.M. Grant, S.M. Williams, Bone and body composition of children and  
524 adolescents with repeated forearm fractures, *J. Bone Miner. Res.* 20 (2005) 2090–2096.  
525 doi:10.1359/JBMR.050820.
- 526 [11] K. Hind, M. Burrows, Weight-bearing exercise and bone mineral accrual in children and  
527 adolescents: A review of controlled trials, *Bone.* 40 (2007) 14–27.  
528 doi:10.1016/j.bone.2006.07.006.

- 529 [12] H.M. Frost, Bone “mass” and the “mechanostat”: a proposal., *Anat. Rec.* 219 (1987) 1–9.  
530 doi:10.1002/ar.1092190104.
- 531 [13] P. Dimitri, J.K. Wales, N. Bishop, Adipokines, bone-derived factors and bone turnover in  
532 obese children; evidence for altered fat-bone signalling resulting in reduced bone mass, *Bone*.  
533 48 (2011) 189–196. doi:10.1016/j.bone.2010.09.034.
- 534 [14] M. Kawai, F.J.A. de Paula, C.J. Rosen, New insights into osteoporosis: The bone-fat  
535 connection, *J. Intern. Med.* 272 (2012) 317–329. doi:10.1111/j.1365-2796.2012.02564.x.
- 536 [15] I.R. Reid, Fat and bone., *Arch. Biochem. Biophys.* 503 (2010) 20–7.  
537 doi:10.1016/j.abb.2010.06.027.
- 538 [16] M.F. Holick, Vitamin D Deficiency, *N. Engl. J. Med.* 357 (2007) 266–281.  
539 doi:10.1056/NEJMr070553.
- 540 [17] C.E. Moore, Y. Liu, Low serum 25-hydroxyvitamin D concentrations are associated with total  
541 adiposity of children in the United States: National Health and Examination Survey 2005 to  
542 2006., *Nutr. Res.* 36 (2016) 72–9. doi:10.1016/j.nutres.2015.11.003.
- 543 [18] M. Brotto, L. Bonewald, Bone and muscle: Interactions beyond mechanical., *Bone*. 80 (2015)  
544 109–14. doi:10.1016/j.bone.2015.02.010.
- 545 [19] A.D. Bakker, R.T. Jaspers, IL-6 and IGF-1 Signaling Within and Between Muscle and Bone:  
546 How Important is the mTOR Pathway for Bone Metabolism?, *Curr. Osteoporos. Rep.* 13  
547 (2015) 131–9. doi:10.1007/s11914-015-0264-1.
- 548 [20] L. Xu, Q. Wang, Q. Wang, A. Lyytikäinen, T. Mikkola, E. Völgyi, S. Cheng, P. Wiklund, E.  
549 Munukka, P. Nicholson, M. Alén, S. Cheng, Concerted actions of insulin-like growth factor 1,  
550 testosterone, and estradiol on peripubertal bone growth: a 7-year longitudinal study., *J. Bone*  
551 *Miner. Res.* 26 (2011) 2204–11. doi:10.1002/jbmr.422.
- 552 [21] T.A.L. Wren, H.J. Kalkwarf, B.S. Zemel, J.M. Lappe, S. Oberfield, J.A. Shepherd, K.K.  
553 Winer, V. Gilsanz, Longitudinal tracking of dual-energy X-ray absorptiometry bone measures  
554 over 6 years in children and adolescents: Persistence of low bone mass to maturity, *J. Pediatr.*  
555 164 (2014) 1280–1285.e2. doi:10.1016/j.jpeds.2013.12.040.
- 556 [22] NIH Consensus Development Panel on Osteoporosis Prevention, Diagnosis, and Therapy.  
557 Osteoporosis Prevention, Diagnosis, and Therapy, *JAMA J. Am. Med. Assoc.* 285 (2001) 785–  
558 795. doi:10.1001/jama.285.6.785.
- 559 [23] M. Ng, T. Fleming, M. Robinson, B. Thomson, N. Graetz, C. Margono, E.C. Mullany, S.  
560 Biryukov, C. Abbafati, S.F. Abera, J.P. Abraham, N.M.E. Abu-Rmeileh, T. Achoki, F.S.  
561 AlBuhairan, Z.A. Alemu, R. Alfonso, M.K. Ali, R. Ali, N.A. Guzman, W. Ammar, P. Anwari,  
562 A. Banerjee, S. Barquera, S. Basu, D.A. Bennett, Z. Bhutta, J. Blore, N. Cabral, I.C. Nonato,

563 J.-C. Chang, R. Chowdhury, K.J. Courville, M.H. Criqui, D.K. Cundiff, K.C. Dabhadkar, L.  
564 Dandona, A. Davis, A. Dayama, S.D. Dharmaratne, E.L. Ding, A.M. Durrani, A. Esteghamati,  
565 F. Farzadfar, D.F.J. Fay, V.L. Feigin, A. Flaxman, M.H. Forouzanfar, A. Goto, M.A. Green,  
566 R. Gupta, N. Hafezi-Nejad, G.J. Hankey, H.C. Harewood, R. Havmoeller, S. Hay, L.  
567 Hernandez, A. Husseini, B.T. Idrisov, N. Ikeda, F. Islami, E. Jahangir, S.K. Jassal, S.H. Jee,  
568 M. Jeffreys, J.B. Jonas, E.K. Kabagambe, S.E.A.H. Khalifa, A.P. Kengne, Y.S. Khader, Y.-H.  
569 Khang, D. Kim, R.W. Kimokoti, J.M. Kinge, Y. Kokubo, S. Kosen, G. Kwan, T. Lai, M.  
570 Leinsalu, Y. Li, X. Liang, S. Liu, G. Logroscino, P.A. Lotufo, Y. Lu, J. Ma, N.K. Mainoo,  
571 G.A. Mensah, T.R. Merriman, A.H. Mokdad, J. Moschandreas, M. Naghavi, A. Naheed, D.  
572 Nand, K.M.V. Narayan, E.L. Nelson, M.L. Neuhauser, M.I. Nisar, T. Ohkubo, S.O. Oti, A.  
573 Pedroza, D. Prabhakaran, N. Roy, U. Sampson, H. Seo, S.G. Sepanlou, K. Shibuya, R. Shiri,  
574 I. Shiue, G.M. Singh, J.A. Singh, V. Skirbekk, N.J.C. Stapelberg, L. Sturua, B.L. Sykes, M.  
575 Tobias, B.X. Tran, L. Trasande, H. Toyoshima, S. van de Vijver, T.J. Vasankari, J.L. Veerman,  
576 G. Velasquez-Melendez, V.V. Vlassov, S.E. Vollset, T. Vos, C. Wang, X. Wang, E.  
577 Weiderpass, A. Werdecker, J.L. Wright, Y.C. Yang, H. Yatsuya, J. Yoon, S.-J. Yoon, Y. Zhao,  
578 M. Zhou, S. Zhu, A.D. Lopez, C.J.L. Murray, E. Gakidou, Global, regional, and national  
579 prevalence of overweight and obesity in children and adults during 1980–2013: a systematic  
580 analysis for the Global Burden of Disease Study 2013, *Lancet*. 384 (2014) 766–781.  
581 doi:10.1016/S0140-6736(14)60460-8.

582 [24] A. Saari, U. Sankilampi, M.-L. Hannila, V. Kiviniemi, K. Kesseli, L. Dunkel, New Finnish  
583 growth references for children and adolescents aged 0 to 20 years: Length/height-for-age,  
584 weight-for-length/height, and body mass index-for-age., *Ann. Med.* 43 (2011) 235–248.  
585 doi:10.3109/07853890.2010.515603.

586 [25] S. Soininen, A.-M. Eloranta, V. Lindi, T. Venäläinen, N. Zaproudina, A. Mahonen, T.A.  
587 Lakka, Determinants of serum 25-hydroxyvitamin D concentration in Finnish children: the  
588 Physical Activity and Nutrition in Children (PANIC) study, *Br. J. Nutr.* 25 (2016) 1–12.  
589 doi:10.1017/S0007114515005292.

590 [26] S. Soininen, A.-M. Eloranta, V. Lindi, T.A. Lakka, Response: food fortification as a means to  
591 increase vitamin D intake., *Br. J. Nutr.* 116 (2016) 1–2. doi:10.1017/S0007114516002890.

592 [27] A. Mäntyselkä, J. Jääskeläinen, V. Lindi, A. Viitasalo, T. Tompuri, R. Voutilainen, T.A.  
593 Lakka, The presentation of adrenarche is sexually dimorphic and modified by body adiposity.,  
594 *J. Clin. Endocrinol. Metab.* 99 (2014) 3889–94. doi:10.1210/jc.2014-2049.

595 [28] J. Kratzsch, A. Lammert, A. Bottner, B. Seidel, G. Mueller, J. Thiery, J. Hebebrand, W. Kiess,  
596 Circulating soluble leptin receptor and free leptin index during childhood, puberty, and

adolescence., *J. Clin. Endocrinol. Metab.* 87 (2002) 4587–94. doi:10.1210/jc.2002-020001.

[29] P. Utriainen, R. Voutilainen, J. Jaaskelainen, Continuum of phenotypes and sympathoadrenal function in premature adrenarche, *Eur. J. Endocrinol.* 160 (2009) 657–665. doi:10.1530/EJE-08-0367.

[30] U. Sankilampi, M.-L. Hannila, A. Saari, M. Gissler, L. Dunkel, New population-based references for birth weight, length, and head circumference in singletons and twins from 23 to 43 gestation weeks., *Ann. Med.* 45 (2013) 446–54. doi:10.3109/07853890.2013.803739.

[31] V. Singhal, E.A. Lawson, K.E. Ackerman, P.K. Fazeli, H. Clarke, H. Lee, K. Eddy, D.A. Marengi, N.P. Derrico, M.L. Bouxsein, M. Misra, Irisin levels are lower in young amenorrheic athletes compared with eumenorrheic athletes and non-athletes and are associated with bone density and strength estimates., *PLoS One.* 9 (2014) e100218. doi:10.1371/journal.pone.0100218.

[32] G. Colaianni, C. Cuscito, T. Mongelli, P. Pignataro, C. Buccoliero, P. Liu, P. Lu, L. Sartini, M. Di Comite, G. Mori, A. Di Benedetto, G. Brunetti, T. Yuen, L. Sun, J.E. Reseland, S. Colucci, M.I. New, M. Zaidi, S. Cinti, M. Grano, The myokine irisin increases cortical bone mass., *Proc. Natl. Acad. Sci. U. S. A.* 112 (2015) 12157–62. doi:10.1073/pnas.1516622112.

[33] A. Palermo, R. Strollo, E. Maddaloni, D. Tuccinardi, L. D’Onofrio, S.I. Briganti, G. Defeudis, M. De Pascalis, M.C. Lazzaro, G. Colleluori, S. Manfrini, P. Pozzilli, N. Napoli, Irisin is associated with osteoporotic fractures independently of bone mineral density, body composition or daily physical activity., *Clin. Endocrinol. (Oxf).* 82 (2015) 615–9. doi:10.1111/cen.12672.

[34] J.M. Kindler, N.K. Pollock, E.M. Laing, N.T. Jenkins, A. Oshri, C. Isales, M. Hamrick, R.D. Lewis, Insulin Resistance Negatively Influences the Muscle-Dependent IGF-1-Bone Mass Relationship in Premenarcheal Girls, *J. Clin. Endocrinol. Metab.* 101 (2016) 199–205. doi:10.1210/jc.2015-3451.

[35] J.M. Kindler, N.K. Pollock, E.M. Laing, A. Oshri, N.T. Jenkins, C.M. Isales, M.W. Hamrick, K.-H. Ding, D.B. Hausman, G.P. McCabe, B.R. Martin, K.M. Hill Gallant, S.J. Warden, C.M. Weaver, M. Peacock, R.D. Lewis, Insulin Resistance and the IGF-I-Cortical Bone Relationship in Children Ages 9-13 Years, *J. Bone Miner. Res.* (2017). doi:10.1002/jbmr.3132.

[36] E.M. Clark, A.R. Ness, J.H. Tobias, Adipose tissue stimulates bone growth in prepubertal children., *J. Clin. Endocrinol. Metab.* 91 (2006) 2534–41. doi:10.1210/jc.2006-0332.

[37] M.B. Leonard, J. Shults, B.A. Wilson, A.M. Tershakovec, B.S. Zemel, Obesity during childhood and adolescence augments bone mass and bone dimensions., *Am. J. Clin. Nutr.* 80 (2004) 514–523.

- 631 [38] Z.A. Cole, N.C. Harvey, M. Kim, G. Ntani, S.M. Robinson, H.M. Inskip, K.M. Godfrey, C.  
632 Cooper, E.M. Dennison, Increased fat mass is associated with increased bone size but reduced  
633 volumetric density in pre pubertal children, *Bone*. 50 (2012) 562–567.  
634 doi:10.1016/j.bone.2011.05.005.
- 635 [39] H.T. Viljakainen, M. Pekkinen, E. Saarnio, H. Karp, C. Lamberg-Allardt, O. Makitie, Dual  
636 effect of adipose tissue on bone health during growth, *Bone*. 48 (2011) 212–217.  
637 doi:10.1016/j.bone.2010.09.022.
- 638 [40] A. Viitasalo, D.E. Laaksonen, V. Lindi, A.-M. Eloranta, J. Jääskeläinen, T. Tompuri, S.  
639 Väisänen, H.-M. Lakka, T.A. Lakka, Clustering of Metabolic Risk Factors Is Associated with  
640 High-Normal Levels of Liver Enzymes Among 6- to 8-Year-Old Children: The PANIC Study,  
641 *Metab. Syndr. Relat. Disord.* 10 (2012) 337–343. doi:10.1089/met.2012.0015.
- 642 [41] J. Upadhyay, O.M. Farr, C.S. Mantzoros, The role of leptin in regulating bone metabolism.,  
643 *Metabolism*. 64 (2015) 105–13. doi:10.1016/j.metabol.2014.10.021.
- 644 [42] K. Liu, P. Liu, R. Liu, X. Wu, M. Cai, Relationship between serum leptin levels and bone  
645 mineral density: A systematic review and meta-analysis, *Clin. Chim. Acta*. 444 (2015) 260–  
646 263. doi:10.1016/j.cca.2015.02.040.
- 647 [43] W.L. do Prado, A. de Piano, M. Lazaretti-Castro, M.T. de Mello, S.G. Stella, S. Tufik, C.M.O.  
648 do Nascimento, L.M. Oyama, M.C. Lofrano, L. Tock, D.A. Caranti, A.R. Dâmaso,  
649 Relationship between bone mineral density, leptin and insulin concentration in Brazilian obese  
650 adolescents, *J. Bone Miner. Metab.* 27 (2009) 613–619. doi:10.1007/s00774-009-0082-6.
- 651 [44] A. Sayers, N.J. Timpson, N. Sattar, J. Deanfield, A.D. Hingorani, G. Davey-Smith, J.H.  
652 Tobias, Adiponectin and its association with bone mass accrual in childhood., *J. Bone Miner.*  
653 *Res.* 25 (2010) 2212–20. doi:10.1002/jbmr.116.
- 654 [45] A. Böttner, J. Kratzsch, G. Müller, T.M. Kapellen, S. Blüher, E. Keller, M. Blüher, W. Kiess,  
655 Gender differences of adiponectin levels develop during the progression of puberty and are  
656 related to serum androgen levels., *J. Clin. Endocrinol. Metab.* 89 (2004) 4053–61.  
657 doi:10.1210/jc.2004-0303.
- 658 [46] R. Weiss, J. Dziura, T.S. Burgert, W.V. Tamborlane, S.E. Taksali, C.W. Yeckel, K. Allen, M.  
659 Lopes, M. Savoye, J. Morrison, R.S. Sherwin, S. Caprio, Obesity and the metabolic syndrome  
660 in children and adolescents, *N. Engl. J. Med.* 350 (2004). doi:10.1056/NEJMoa031049.
- 661 [47] N.K. Pollock, P.J. Bernard, K. Wenger, S. Misra, B.A. Gower, J.D. Allison, H. Zhu, C.L.  
662 Davis, Lower bone mass in prepubertal overweight children with prediabetes., *J. Bone Miner.*  
663 *Res.* 25 (2010) 2760–9. doi:10.1002/jbmr.184.
- 664 [48] N.K. Pollock, P.J. Bernard, B. Gutin, C.L. Davis, H. Zhu, Y. Dong, Adolescent obesity, bone

mass, and cardiometabolic risk factors., *J. Pediatr.* 158 (2011) 727–34. doi:10.1016/j.jpeds.2010.11.052.

[49] D.A. Lawlor, N. Sattar, A. Sayers, J.H. Tobias, The association of fasting insulin, glucose, and lipids with bone mass in adolescents: findings from a cross-sectional study., *J. Clin. Endocrinol. Metab.* 97 (2012) 2068–76. doi:10.1210/jc.2011-2721.

[50] R. Lucas, E. Ramos, A. Oliveira, T. Monjardino, H. Barros, Low-grade systemic inflammation and suboptimal bone mineral density throughout adolescence: a prospective study in girls., *Clin. Endocrinol. (Oxf).* 77 (2012) 665–71. doi:10.1111/j.1365-2265.2012.04430.x.

[51] C.S. Tam, K. Clément, L.A. Baur, J. Tordjman, Obesity and low-grade inflammation: a paediatric perspective, *Obes. Rev.* 11 (2010) 118–126. doi:10.1111/j.1467-789X.2009.00674.x.

[52] H.A. Bischoff-Ferrari, E. Giovannucci, W.C. Willett, T. Dietrich, B. Dawson-Hughes, Estimation of optimal serum concentrations of 25-hydroxyvitamin D for multiple health outcomes, *Am. J. Clin. Nutr.* 84 (2006) 18–28.

[53] Dietary Reference Intakes for Calcium and Vitamin D, National Academies Press, Washington, D.C., 2011. doi:10.17226/13050.

[54] C. Lamberg-Allardt, M. Brustad, H.E. Meyer, L. Steingrimsdottir, Vitamin D - a systematic literature review for the 5th edition of the Nordic Nutrition Recommendations., *Food Nutr. Res.* 57 (2013) 1–31. doi:10.3402/fnr.v57i0.22671.

[55] M.F. Holick, N.C. Binkley, H.A. Bischoff-Ferrari, C.M. Gordon, D.A. Hanley, R.P. Heaney, M.H. Murad, C.M. Weaver, Evaluation, treatment, and prevention of vitamin D deficiency: An endocrine society clinical practice guideline, *J. Clin. Endocrinol. Metab.* 96 (2011) 1911–1930. doi:10.1210/jc.2011-0385.

[56] C. Braegger, C. Campoy, V. Colomb, T. Decsi, M. Domellöf, M. Fewtrell, I. Hojsak, W. Mihatsch, C. Molgaard, R. Shamir, D. Turck, J. van Goudoever, Vitamin D in the healthy European paediatric population., *J. Pediatr. Gastroenterol. Nutr.* 56 (2013) 692–701. doi:10.1097/MPG.0b013e31828f3c05.

[57] P. Arundel, S.F. Ahmed, J. Allgrove, N.J. Bishop, C.P. Burren, B. Jacobs, M.Z. Mughal, A.C. Offiah, N.J. Shaw, British Paediatric and Adolescent Bone Group’s position statement on vitamin D deficiency, *BMJ.* 345 (2012) e8182–e8182. doi:10.1136/bmj.e8182.

[58] S. Vandewalle, Y. Taes, T. Fiers, K. Toye, E. Van Caenegem, J.-M. Kaufman, J. De Schepper, Relation of adrenal-derived steroids with bone maturation, mineral density and geometry in healthy prepubertal and early pubertal boys., *Bone.* 69 (2014) 39–46. doi:10.1016/j.bone.2014.09.002.

- 699 [59] A.B. Sopher, A.M. Jean, S.K. Zwany, D.M. Winston, C.B. Pomeranz, J.J. Bell, D.J. McMahon,  
700 A. Hassoun, I. Fennoy, S.E. Oberfield, Bone age advancement in prepubertal children with  
701 obesity and premature adrenarche: possible potentiating factors., *Obesity (Silver Spring)*. 19  
702 (2011) 1259–64. doi:10.1038/oby.2010.305.
- 703 [60] P. Utriainen, J. Jääskeläinen, A. Saarinen, E. Vanninen, O. Mäkitie, R. Voutilainen, Body  
704 composition and bone mineral density in children with premature adrenarche and the  
705 association of LRP5 gene polymorphisms with bone mineral density., *J. Clin. Endocrinol.*  
706 *Metab.* 94 (2009) 4144–51. doi:10.1210/jc.2009-0315.
- 707 [61] V. Saraff, W. Höglér, Osteoporosis in children: Diagnosis and management, *Eur. J.*  
708 *Endocrinol.* 173 (2015) R185–R197. doi:10.1530/EJE-14-0865.
- 709 [62] V.H. Sidoroff, M.K. Ylinen, L.M. Kröger, H.P.J. Kröger, M.O. Korppi, Inhaled corticosteroids  
710 and bone mineral density at school age: a follow-up study after early childhood wheezing.,  
711 *Pediatr. Pulmonol.* 50 (2015) 1–7. doi:10.1002/ppul.22968.
- 712 [63] Y.K. Loke, D. Gilbert, M. Thavarajah, P. Blanco, A.M. Wilson, Bone mineral density and  
713 fracture risk with long-term use of inhaled corticosteroids in patients with asthma: systematic  
714 review and meta-analysis., *BMJ Open*. 5 (2015) e008554. doi:10.1136/bmjopen-2015-008554.
- 715 [64] L. Margulies, M. Horlick, J.C. Thornton, J. Wang, E. Ioannidou, S.B. Heymsfield,  
716 Reproducibility of pediatric whole body bone and body composition measures by dual-energy  
717 X-ray absorptiometry using the GE Lunar Prodigy, *J Clin.Densitom.* 8 (2005) 298–304.  
718 doi:JCD:8:3:298 [pii].
- 719 [65] J.A. Shepherd, L. Wang, B. Fan, V. Gilsanz, H.J. Kalkwarf, J. Lappe, Y. Lu, T. Hangartner,  
720 B.S. Zemel, M. Fredrick, S. Oberfield, K.K. Winer, Optimal monitoring time interval between  
721 DXA measures in children, *J. Bone Miner. Res.* 26 (2011) 2745–2752. doi:10.1002/jbmr.473.
- 722 [66] N.J. Crabtree, A. Arabi, L.K. Bachrach, M. Fewtrell, G. El-Hajj Fuleihan, H.H. Kecs kemethy,  
723 M. Jaworski, C.M. Gordon, Dual-energy x-ray absorptiometry interpretation and reporting in  
724 children and adolescents: The revised 2013 ISCD pediatric official positions, *J. Clin.*  
725 *Densitom.* 17 (2014) 225–242. doi:10.1016/j.jocd.2014.01.003.



726 **Table 1. Characteristics of children.**

	All (n=472)	Girls (n=227)	Boys (n=245)	P-value
Age, y	7.6 (0.4)	7.6 (0.4)	7.6 (0.4)	0.169
Birth weight SDS	-0.05 (1.00)	-0.01 (0.99)	-0.09 (1.01)	0.372
Weight, kg	26.7 (4.9)	26.2 (4.8)	27.3 (4.9)	<b>0.017</b>
Height, cm	128.6 (5.6)	127.7 (5.6)	129.5 (5.4)	<b>0.001</b>
BMI-SDS	-0.20 (1.07)	-0.23 (1.02)	-0.17 (1.11)	0.511
Waist circumference, cm	56.5 (5.7)	55.5 (5.4)	57.5 (5.7)	<b>&lt;0.001</b>
Lean body mass, kg	17.7 (2.2)	16.8 (2.0)	18.6 (2.1)	<b>&lt;0.001</b>
Body fat mass, kg	5.2 (3.3)	5.6 (3.2)	4.7 (3.3)	<b>0.002</b>
Body fat percentage	20.6 (8.5)	23.2 (7.8)	18.2 (8.5)	<b>&lt;0.001</b>
BMD, total body excluding the head, g/cm <sup>2</sup>	0.72 (0.05)	0.72 (0.05)	0.72 (0.05)	0.094
25(OH)D, nmol/l	67.8 (22.7)	65.6 (18.8)	69.8 (25.7)	0.056
DHEAS, µmol/l	0.57 (0.33-0.85)	0.57 (0.33-0.84)	0.58 (0.32-0.85)	0.998
IGF-1, nmol/l	23.1 (7.6)	24.4 (7.3)	22.0 (7.5)	<b>0.001</b>
Insulin, mU/l	4.50 (2.44)	4.74 (2.25)	4.29 (2.58)	<b>0.049</b>
HOMA-IR	0.98 (0.56)	1.01 (0.51)	0.94 (0.59)	0.196
Adiponectin, µg/ml	8.91 (4.09)	8.84 (3.77)	8.97 (4.36)	0.740
Leptin, ng/ml	3.70 (2.70-5.85)	4.30 (3.20-6.80)	3.20 (2.40-4.70)	<b>&lt;0.001</b>
Leptin receptor, ng/ml	40.8 (10.5)	38.6 (9.7)	42.9 (10.9)	<b>&lt;0.001</b>
Free leptin index	9.0 (5.9-16.2)	10.9 (7.5-20.6)	6.9 (5.1-12.6)	<b>&lt;0.001</b>
Irisin, ng/ml	151.5 (53.0)	151.3 (43.3)	151.8 (60.5)	0.918
IL-6, pg/ml	0.90 (0.63-1.47)	0.83 (0.60-1.27)	1.00 (0.63-1.57)	<b>0.016</b>
TNF-α, pg/ml	14.3 (5.2-34.2)	12.7 (5.0-30.4)	15.7 (5.4-37.4)	0.177
hs-CRP, mg/l	0.29 (0.29-0.54)	0.29 (0.29-0.59)	0.29 (0.29-0.49)	0.098

727  
728 The values are presented as mean (SD) for normally distributed variables and median (IQR) for skewed  
729 variables. Differences between girls and boys were tested with independent samples t test for normally  
730 distributed variables and Mann–Whitney’s U test for skewed variables. P-value for differences between girls  
731 and boys.

732  
733 Abbreviations: SDS, standard deviation score; BMI-SDS, body mass index standard deviation score; BMD,  
734 bone mineral density; 25(OH)D, 25-hydroxyvitamin D; DHEAS, dehydroepiandrosterone sulphate; IGF-1,  
735 insulin-like growth factor 1; HOMA-IR: homeostatic model assessment for insulin resistance; adiponectin,  
736 high-molecular weight adiponectin; IL-6, interleukin 6; TNF-α, tumor necrosis factor α; hs-CRP, high-  
737 sensitivity C-reactive protein (values over 10 excluded)

738  
739 Number of children (n) varies from 417 to 472 in different variables; n=472, 227 girls and 245 boys: age,  
740 weight, height, BMI-SDS, waist, lean body mass, body fat mass, BMD; n=463, 222 girls and 241 boys: birth  
741 weight SDS; n=417, 198 girls and 219 boys: 25(OH)D; n= 440, 211 girls and 229 boys DHEAS, IGF-1; n=  
742 456, 216 girls and 240 boys :insulin; n= 452, 215 girls and 237 boys: HOMA-IR; n= 452, 214 girls and 238  
743 boys: adiponectin, leptin, leptin receptor; n= 433, 205 girls and 228 boys: irisin; n= 448, 210 girls and 238  
744 boys: IL-6; n= 450, 211 girls and 239 boys: TNF-α; n= 456, 217 girls and 239 boys: hs-CRP (values over 10  
745 excluded).

746 **Table 2. Determinants of bone mineral density (total body excluding the head) in all children.**

	Model 1		Model 2		Model 3		Model 4	
	Beta	p-value	Beta	p-value	Beta	p-value	Beta	p-value
Lean body mass, kg	0.729	<b>&lt;0.001</b>	0.708	<b>&lt;0.001</b>			0.562	<b>&lt;0.001</b>
Body fat mass, kg	0.594	<b>&lt;0.001</b>	0.358	<b>&lt;0.001</b>	0.365	<b>&lt;0.001</b>		
Birth weight SDS	0.169	<b>&lt;0.001</b>	-0.011	0.786	-0.059	0.103	0.009	0.807
25(OH)D, nmol/l	0.097	<b>0.044</b>	0.086	<b>0.036</b>	0.067	0.075	0.087	<b>0.017</b>
DHEAS, µmol/l	0.178	<b>&lt;0.001</b>	0.100	<b>0.012</b>	0.071	0.052	0.065	0.068
IGF-1, nmol/l	0.188	<b>&lt;0.001</b>	0.037	0.375	0.007	0.844	-0.041	0.275
Insulin, mU/l	0.218	<b>&lt;0.001</b>	0.102	<b>0.010</b>	0.071	<b>0.048</b>	-0.043	0.260
HOMA-IR	0.212	<b>&lt;0.001</b>	0.087	<b>0.028</b>	0.062	0.087	-0.054	0.153
Adiponectin, µg/ml	-0.091	<b>0.049</b>	-0.067	0.082	-0.042	0.232	-0.052	0.126
Leptin, ng/ml	0.397	<b>&lt;0.001</b>	0.275	<b>&lt;0.001</b>	0.245	<b>&lt;0.001</b>	-0.114	0.058
Leptin receptor, ng/ml	-0.432	<b>&lt;0.001</b>	-0.260	<b>&lt;0.001</b>	-0.192	<b>&lt;0.001</b>	0.061	0.168
Free leptin index	0.400	<b>&lt;0.001</b>	0.284	<b>&lt;0.001</b>	0.245	<b>&lt;0.001</b>	0.012	0.825
Irisin, ng/ml	0.105	<b>0.026</b>	0.079	<b>0.048</b>	0.072	<b>0.047</b>	0.075	<b>0.034</b>
IL-6, pg/l	0.042	0.370	0.031	0.422	0.007	0.836	-0.001	0.982
TNF-α pg/ml	0.009	0.844	0.022	0.568	0.014	0.679	0.032	0.341
hs-CRP, mg/l	0.111	<b>0.014</b>	0.088	<b>0.023</b>	0.098	<b>0.005</b>	0.003	0.941

747

748 The values are standardized regression coefficients (β) and p-values from linear regression models.

749

750 Model 1: Each variable was entered in linear regression analysis adjusted for age and sex.

751 Model 2: Each variable was entered in linear regression analysis adjusted for age, sex, and height.

752 Model 3: Each variable was entered in linear regression analysis adjusted for age, sex, height, and lean mass.

753 Model 4: Each variable was entered in linear regression analysis adjusted for age, sex, height, and fat mass.

754

755 Abbreviations: SDS, standard deviation score; BMI-SDS, body mass index standard deviation score; BMD, bone mineral density; 25(OH)D, 25-hydroxyvitamin D; DHEAS, dehydroepiandrosterone sulphate; IGF-1, insulin-like growth factor 1; HOMA-IR: homeostatic model assessment for insulin resistance; adiponectin, high-molecular weight adiponectin; IL-6, interleukin 6; TNF-α, tumor necrosis factor α; hs-CRP, high-sensitivity C-reactive protein (values over 10 excluded)

760

761 Number of children (n) varies from 417 to 472 in different variables; n=472: BMD, lean body mass, body fat mass; n=463: birth weight SDS; n=417: 25(OH)D; n= 440: DHEAS, IGF-1; n= 456: insulin; n= 452: HOMA-IR; n= 452: adiponectin, leptin, leptin receptor; n= 433: irisin; n= 448: IL-6; n= 450: TNF-α; n= 456: hs-CRP (values over 10 excluded).

764

765

766 **Table 3. Determinants of bone mineral density (total body excluding the head) in girls.**

	Model 1		Model 2		Model 3		Model 4	
	Beta	p-value	Beta	p-value	Beta	p-value	Beta	p-value
Lean body mass, kg	0.663	<0.001	0.571	<0.001			0.459	<0.001
Body fat mass, kg	0.612	<0.001	0.439	<0.001	0.382	<0.001		
Birth weight SDS	0.187	0.004	-0.009	0.870	-0.047	0.357	-0.016	0.749
25(OH)D, nmol/l	0.097	0.044	0.042	0.464	0.048	0.350	0.035	0.473
DHEAS, µmol/l	0.256	<0.001	0.095	0.095	0.078	0.132	0.030	0.551
IGF-1, nmol/l	0.278	<0.001	0.083	0.157	0.060	0.258	0.040	0.422
Insulin, mU/l	0.231	<0.001	0.139	0.010	0.120	0.015	0.019	0.693
HOMA-IR	0.228	<0.001	0.136	0.012	0.120	0.014	0.015	0.758
Adiponectin, µg/ml	-0.061	0.352	-0.060	0.263	-0.042	0.394	-0.018	0.706
Leptin, ng/ml	0.425	<0.001	0.305	<0.001	0.279	<0.001	-0.064	0.417
Leptin receptor, ng/ml	-0.432	<0.001	-0.271	<0.001	-0.226	<0.001	-0.074	0.198
Free leptin index	0.434	<0.001	0.321	<0.001	0.286	<0.001	0.034	0.629
Irisin, ng/ml	0.175	0.008	0.108	0.052	0.094	0.063	0.073	0.126
IL-6, pg/l	0.120	0.070	0.061	0.269	0.038	0.452	0.021	0.656
TNF-α pg/ml	0.054	0.416	0.065	0.232	0.024	0.630	0.052	0.270
hs-CRP, mg/l	0.070	0.283	0.070	0.198	0.092	0.065	-0.030	0.563

767

768 The values are standardized regression coefficients (β) and p-values from linear regression models.

769

770 Model 1: Each variable was entered in linear regression analysis adjusted for age.

771 Model 2: Each variable was entered in linear regression analysis adjusted for age and height.

772 Model 3: Each variable was entered in linear regression analysis adjusted for age, height, and lean mass.

773 Model 4: Each variable was entered in linear regression analysis adjusted for age, height, and fat mass.

774

775 Abbreviations: SDS, standard deviation score; BMI-SDS, body mass index standard deviation score; BMD, bone mineral density; 25(OH)D, 25-hydroxyvitamin D; DHEAS, dehydroepiandrosterone sulphate; IGF-1, insulin-like growth factor 1; HOMA-IR: homeostatic model assessment for insulin resistance; adiponectin, high-molecular weight adiponectin; IL-6, interleukin 6; TNF-α, tumor necrosis factor α; hs-CRP, high-sensitivity C-reactive protein (\*values over 10 excluded)

780

781 Number of girls (n) varies from 205 to 227 in different variables; n=227: BMD, lean body mass, body fat mass; n=222: birth weight SDS; n=198: 25(OH)D; n=211: DHEAS, IGF-1; n=216: insulin; n=215: HOMA-IR; n=214: adiponectin, leptin, leptin receptor; n=205: irisin; n= 210: IL-6; n=211: TNF-α; n=217: hs-CRP (values over 10 excluded).

784

785 **Table 4. Determinants of bone mineral density (total body excluding the head) in boys.**

	Model 1		Model 2		Model 3		Model 4	
	Beta	p-value	Beta	p-value	Beta	p-value	Beta	p-value
Lean body mass, kg	0.664	< <b>0.001</b>	0.746	< <b>0.001</b>			0.591	< <b>0.001</b>
Body fat mass, kg	0.568	< <b>0.001</b>	0.435	< <b>0.001</b>	0.337	< <b>0.001</b>		
Birth weight SDS	0.145	<b>0.023</b>	-0.029	0.619	-0.074	0.153	0.031	0.555
25(OH)D, nmol/l	0.123	0.070	0.117	<b>0.045</b>	0.079	0.145	0.123	<b>0.023</b>
DHEAS, µmol/l	0.139	<b>0.036</b>	0.108	0.062	0.070	0.182	0.088	0.095
IGF-1, nmol/l	0.094	0.152	-0.007	0.905	-0.045	0.393	-0.108	<b>0.049</b>
Insulin, mU/l	0.211	<b>0.001</b>	0.079	0.172	0.039	0.456	-0.089	0.132
HOMA-IR	0.209	<b>0.001</b>	0.057	0.975	0.028	0.598	-0.108	0.069
Adiponectin, µg/ml	-0.127	0.050	-0.080	0.152	-0.055	0.278	-0.080	0.115
Leptin, ng/ml	0.346	< <b>0.001</b>	0.229	< <b>0.001</b>	0.187	< <b>0.001</b>	-0.266	<b>0.006</b>
Leptin receptor, ng/ml	-0.418	< <b>0.001</b>	-0.249	< <b>0.001</b>	-0.161	<b>0.004</b>	-0.056	0.401
Free leptin index	0.353	< <b>0.001</b>	0.236	< <b>0.001</b>	0.180	< <b>0.001</b>	-0.165	0.063
Irisin, ng/ml	0.056	0.400	0.058	0.312	0.053	0.310	0.073	0.168
IL-6, pg/l	-0.019	0.770	0.006	0.914	-0.023	0.650	0.019	0.992
TNF-α pg/ml	-0.020	0.762	-0.010	0.854	0.015	0.766	0.015	0.768
hs-CRP*, mg/l	0.138	<b>0.032</b>	0.101	0.068	0.083	0.096	0.024	0.650

786

787 The values are standardized regression coefficients (β) and p-values from linear regression models.

788

789 Model 1: Each variable was entered in linear regression analysis adjusted for age.

790 Model 2: Each variable was entered in linear regression analysis adjusted for age and height.

791 Model 3: Each variable was entered in linear regression analysis adjusted for age, height, and lean mass.

792 Model 4: Each variable was entered in linear regression analysis adjusted for age, height, and body fat mass.

793

794 Abbreviations: SDS, standard deviation score; BMI-SDS, body mass index standard deviation score; BMD, bone mineral density; 25(OH)D, 25-hydroxyvitamin D; DHEAS, dehydroepiandrosterone sulphate; IGF-1, insulin-like growth factor 1; HOMA-IR: homeostatic model assessment for insulin resistance; adiponectin, high-molecular weight adiponectin; IL-6, interleukin 6; TNF-α, tumor necrosis factor α; hs-CRP, high-sensitivity C-reactive protein (values over 10 excluded)

799

800 Number of boys (n) varies from 219 to 245 in different variables; n= 245: BMD, lean mass, body fat mass; n=241: birth weight SDS; n=219: 25(OH)D; n=229: DHEAS, IGF-1; n=240: insulin; n=237: HOMA-IR; n=238: adiponectin, leptin, leptin receptor; n=228: irisin; n=238: IL-6; n=239: TNF-α; n=239: hs-CRP (values over 10 excluded).

803