



## Understanding of anesthesia – Why consciousness is essential for life and not based on genes

František Baluška, Ken Yokawa, Stefano Mancuso & Keith Baverstock

To cite this article: František Baluška, Ken Yokawa, Stefano Mancuso & Keith Baverstock (2016) Understanding of anesthesia – Why consciousness is essential for life and not based on genes, Communicative & Integrative Biology, 9:6, e1238118, DOI: [10.1080/19420889.2016.1238118](https://doi.org/10.1080/19420889.2016.1238118)

To link to this article: <https://doi.org/10.1080/19420889.2016.1238118>



© 2016 The Author(s). Published with license by Taylor & Francis. © František Baluška, Ken Yokawa, Stefano Mancuso, and Keith Baverstock.



Published online: 04 Nov 2016.



Submit your article to this journal [↗](#)



Article views: 886



View related articles [↗](#)



View Crossmark data [↗](#)



Citing articles: 4 View citing articles [↗](#)

## Understanding of anesthesia – Why consciousness is essential for life and not based on genes

František Baluška<sup>a</sup>, Ken Yokawa<sup>a</sup>, Stefano Mancuso<sup>b</sup>, and Keith Baverstock<sup>c</sup>

<sup>a</sup>IZMB, University of Bonn, Kirschalle, Bonn, Germany; <sup>b</sup>Department of Plant, Soil and Environmental Science & LINV, University of Florence, Sesto Fiorentino, Italy; <sup>c</sup>Department of Environmental Science, University of Eastern Finland, Kuopio, Finland

### ABSTRACT

Anesthesia and consciousness represent 2 mysteries not only for biology but also for physics and philosophy. Although anesthesia was introduced to medicine more than 160 y ago, our understanding of how it works still remains a mystery. The most prevalent view is that the human brain and its neurons are necessary to impose the effects of anesthetics. However, the fact is that all life can be anesthetized. Numerous theories have been generated trying to explain the major impact of anesthetics on our human-specific consciousness; switching it off so rapidly, but no single theory resolves this enduring mystery. The speed of anesthetic actions precludes any direct involvement of genes. Lipid bilayers, cellular membranes, and critical proteins emerge as the most probable primary targets of anesthetics. Recent findings suggest, rather surprisingly, that physical forces underlie both the anesthetic actions on living organisms as well as on consciousness in general.

### ARTICLE HISTORY

Received 10 July 2015  
Revised 5 August 2016  
Accepted 14 September 2016

### KEYWORDS

anesthesia; anesthetics;  
consciousness; ether;  
ethylene; genes; lipids;  
membranes; plants; xenon

### Introduction

The discovery of human anesthesia in 1846, by William T. Morton, especially its fast and reversible induction by ether; marks new era in our understanding of life.<sup>1-3</sup> This was a unique discovery which rapidly revolutionized medicine, but turned out to be difficult to explain.<sup>1,2</sup> Despite numerous theories proposed our understanding of anesthesia remains obscure.<sup>2,3</sup> Among the many theories trying to explain anesthesia, 3 stand-out as the most influential. Firstly, Claude Bernard and his theory colloidal-coagulation of protoplasm,<sup>2,4</sup> followed by Hans Horst Meyer and Charles Ernest Overton with their “Meyer-Overton rule,” based on lipid solubility of anesthetics, have dominated the field since the late 1960s.<sup>2</sup> In the 1980s, the protein/receptor theory became dominant in explaining anesthesia via proteins acting as specific receptors for anesthetics.<sup>5</sup> However, inconsistencies with this theory called for a further updates.<sup>3,6-10</sup>

### All life can be anesthetized

The father of experimental biology, Claude Bernard, performed numerous experimental studies which allowed him to conclude “that all life is defined by the

susceptibility to anesthesia.”<sup>2,11,12</sup> Claude Bernard’s paradigm is still valid today as all organisms, even prokaryotic bacteria, are sensitive to anesthetics.<sup>8,9,12,13-15</sup> Sub-cellular organelles such as mitochondria and chloroplasts are sensitive to anesthetics as well,<sup>16-21</sup> which is in line with the prokaryotic sensitivity of both membranes and proteins recorded in bacteria.<sup>14,22-24</sup> Finally, several sub-cellular processes based on the cytoskeleton, such as cytoplasmic streaming and phagocytosis, are inhibited by exposure to diverse anesthetics.<sup>6,25-30</sup>

### Early evolutionary origins of anesthesia: Endogenous anesthetics help to cope with stress

As discussed by James Sonner, the variability in response to anesthetics is extremely small in comparison to other drugs.<sup>8,31</sup> Moreover, the wide molecular diversity of compounds acting as anesthetics is very large, and additionally, the mystery of universal sensitivity of all living organisms to these compounds remains. All this suggests, in line with the Claude Bernard thesis, that the ability to respond to anesthetics is essential for life.<sup>8</sup> One possibility proposed recently by James Sonner and Robert Cantor is the existence of endogenous anesthetics which modulate organismal consciousness.<sup>9</sup> In fact, there are several metabolites that induce

**CONTACT** František Baluška  [baluska@uni-bonn.de](mailto:baluska@uni-bonn.de)  IZMB, University of Bonn, Kirschalle 1, 53115 Bonn, Germany; Keith Baverstock  [keith.baverstock@uef.fi](mailto:keith.baverstock@uef.fi)

© 2016 František Baluška, Ken Yokawa, Stefano Mancuso, and Keith Baverstock. Published with license by Taylor & Francis.

This is an Open Access article distributed under the terms of the Creative Commons Attribution-Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>), which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited. The moral rights of the named author(s) have been asserted.

loss of consciousness if available in sufficiently large amounts. Sonner and Cantor discussed some, for example, ammonia,<sup>32</sup> acetone,  $\beta$ -hydroxybutyric acid,<sup>33</sup> and propionic acid.<sup>34</sup> Importantly, protective actions of endogenous anesthetics are active at the lipid bilayer of membranes,<sup>8,9,35</sup> which explains why even bacteria are sensitive to anesthetics. In addition, the protective actions of anesthetics include also cardioprotection,<sup>36-38</sup> protection from retinal damages,<sup>39</sup> and immunoprotection.<sup>40</sup> Finally, due to their actions also on bacteria, anesthetics also have antimicrobial effects.<sup>23,41</sup>

### Microbes, algae and plants release large amounts of volatile anesthetics

It is not generally appreciated that algae and plants release abundantly volatiles including well-known anesthetics such as chloroform, divinyl ether, ethylene, and methyl halides,<sup>42-48</sup> as well as n-alkanols which also have anesthetic actions.<sup>49,50</sup> Plant volatiles are released in such large amounts that they have a strong impact on the Earth's biosphere and atmosphere.<sup>43,47</sup> Importantly, algae and plants release these substances especially if under stress.<sup>44,46,48</sup> In addition, the anesthetic nitrous oxide is released into atmosphere in large amounts from soils and oceans.<sup>1-53</sup> Unfortunately, the authors of these papers fail to mention and discuss the fact that many of these stress-released algal, microbial and plant volatiles are also anesthetics. For example, nitrous oxide is almost exclusively regarded as a greenhouse gas, whereas divinyl ether is discussed typically as oxylipin<sup>54,55</sup> and ethylene as plant stress hormone.<sup>56,57</sup> However, the possibility that stressed organisms produce anesthetics to cope better with their stress should be considered. Besides classical anesthetics, stressed plant synthesize and produce many pain-relieving compounds,<sup>58-62</sup> including ethanol which is produced in plants via the synthetic activity of alcohol dehydrogenase, which acts in reverse in plants.<sup>63,64</sup> In stressed roots, for example, all 3 endogenous anesthetics (ethylene, divinyl ether, ethanol) are co-produced, which suggests delicate control of putative, still hypothetical, plant anesthesia. Relevant in this respect is the fact that all these 3 endogenous anesthetics are co-produced also in plant fruits, which flowering plants have evolved to be eaten alive by diverse animals. Moreover, ethylene, ethanol and other anesthetics also act to break the dormancy in plant seeds.<sup>65</sup>

### Surprising status of ethylene: Ancient endogenous anesthetic essential for life?

Ethylene, a hydrocarbon and the simplest alkene, is a colourless flammable gas, which is widely used in the

chemical industry. In biology, it is most famously known as a plant stress hormone. Less known is that ethylene is a very potent general anesthetic, which was used in human surgery as it has minimum side-effects and recovery from anesthesia is very rapid.<sup>66-68</sup> Ethylene has similar physical and lipid solubility properties to xenon,<sup>66</sup> and both these anesthetics have the least side effects. Importantly, not only plants, but also bacteria, fungi, algae and lichens are known to produce ethylene when under stress.<sup>69-80</sup> This suggests that ethylene is a molecule with fundamental relevance for life. On the basis of the anesthetic properties of ethylene and ether, Chauncey Leake synthesized divinyl ether, and showed that it had excellent anesthetic properties.<sup>81,82</sup> Divinyl ether maintains all the positive properties of ethylene, but is also more potent.<sup>83</sup> Intriguingly, stressed plants produce endogenously both ethylene and divinyl ether.<sup>84-87</sup> It is becoming obvious that plants, and their endogenous anesthetics, will turn out to be highly relevant for our understanding of the evolutionary origins of anesthesia. Very relevant in this respect are local anesthetics most of which are derived from the plant alkaloid cocaine.<sup>88,89</sup> Interestingly, there are also other plant alkaloids with properties of local anesthetics; such as atropine,<sup>90</sup> menthol,<sup>91,92</sup> and several other alkaloids.<sup>93,94</sup>

### The relevance of endogenous controls over anesthesia for behavior and survival

A coherent and robust concept is emerging from the data discussed in this paper, which suggests that endogenous anesthetics are essential for the survival of plants, allowing them to cope with stress, to enter and break dormancy, as well as to generate tasteful fruits 'designed' to be eaten alive by animals and humans (for the sake of effective reproduction of flowering plants). Similarly, bacteria and fungi generate ethylene under stressful situations and there are several indications of endogenous anesthetics in animals and humans. In metabolic human diseases, several metabolites with anesthetic features accumulate in such amounts that they can impose temporary or permanent loss of consciousness (reviewed in ref. 9). Relevant in this respect is the well-known phenomena of transient loss of consciousness (sometimes referred as syncope, fainting, or blackout) which can be induced by diverse stress situations, serious wounding, as well as by powerful emotional stresses.<sup>95-99</sup> Transient loss of consciousness apparently has relevance to survival. It was proposed that this phenomenon has evolved in ancient times as an effective defense mechanism,<sup>97,99</sup> providing, in addition, protection against sensory overload. Moreover, it might also be the case that some of the numerous examples of so-called apparent death


(*thanatopsis*) behavior in predator-threatened animals are not the result of deceptive/ mimicry behavior by the animal, (as typically interpreted) but rather due to the syncope-like transient loss of consciousness. Even at the cellular level, both stress and anesthetics can have similar paralyzing outcomes.<sup>100-102</sup>

Since ancient times, humans have been using natural anesthetics produced by plants and fungi to impose anesthesia, induce altered sensory states as well as psychedelic experiences.<sup>60,61,63-106</sup> In fact, human evolution is well known to be shaped by the consumption of alcohol.<sup>107-109</sup> This feature is shared not only with apes<sup>110</sup> but also with other animals such as wild treeshrews,<sup>111</sup> and even insects, which can also develop alcoholism under stress challenges.<sup>112-114</sup>

Furthermore, there are several similarities between the deep phases of sleep (REM sleep) and the state of anesthesia.<sup>115-117</sup> Although sleep and anesthesia are different phenomena, the underlying neuronal processes are common for both forms of the loss of consciousness.<sup>117-119</sup> Although the precise roles of sleep and anesthesia are not fully understood yet, it is clear that sleep is essential for cognitive and survival reasons.<sup>120-122</sup> For example, REM sleep was proposed to generate a proto-conscious states relevant for the formation of the full-blown waking consciousness.<sup>123</sup> Similarly sleep and anesthesia are under endogenous control,<sup>124</sup> suggesting that the state of anesthesia has important, albeit still elusive, functions for organisms. Importantly, the sleep/ waking cycle controls sensitivity to anesthetics in *Drosophila*.<sup>125</sup> Relevant in this respect are also numerous examples where pain perceptions have been shown to be affected by expectations, cognition and meditation.<sup>126-128</sup> Moreover, pain responses in humans are mediated not only via conscious but also via unconscious processes.<sup>129-131</sup> Finally, there are close similarities between anesthesia and coma.<sup>132</sup>

### Genes are not involved in switching off/on of consciousness via anesthetics

Anesthetics provided in the appropriate concentrations switch-off human-specific consciousness within a few seconds, precluding any role whatsoever for DNA and gene expression in those actions (Fig. 1). If anesthetics are maintained at their active levels, loss of consciousness is permanent as long as the anesthetics are present at critical levels. After their removal, the recovery of consciousness is often very rapid. Although some anesthetics have side effects, and can even be toxic (e.g. chloroform), the most effective ones – for example xenon, ethylene, and vinyl ether allow fast and smooth recovery. Again, the speed of regaining the consciousness precludes any



**Figure 1.** Fast loss and gain of consciousness after exposure and removal of anesthetics is based on primary processes linked to the plasma membrane (ion fluxes, electric activities, endocytic vesicle recycling), whereas changes in gene expression are playing only secondary roles.


active role for gene expression also in this process (Fig. 1). Of course, genes are relevant for the sensitivity of organisms to anesthetics and several mutants have been characterized which are less sensitive, or even not sensitive at all, to some of the anesthetics. But this is just due to the modifications or absence of critical proteins and membranes (Fig. 1) which induce the anesthetic action. The absence of gene involvement should not necessarily be surprising. Human erythrocytes, which have no nucleus and therefore no genes, still undergo the complex cellular process of circadian rhythm.<sup>133</sup> Genes may act to adjust circadian rhythm for variation in the regularity of the day/ night cycle, as, for example, incurred by traveling to other time zones, but they are not causal in circadian rhythm itself. This is clearly demonstrated by incubating 3 enzymes extracted from cyanobacteria with ATP: a relatively temperature independent 24 hour cycle of phosphorylation of one of the enzymes is observed.<sup>134</sup>

### Why are neurons exquisitely sensitive to anesthetics: clues from plants and chloroplasts?

There are several mysteries associated with anesthetics and anesthesia. One of them is the fact that neurons are more sensitive to anesthetics compared to other cells. Claude Bernard was the first to realize that depending upon concentration of an anesthetic, there are several stages of anesthesia. The first is loss of awareness and pain perception, but all vital biochemical processes are unaffected, the second is inhibition of respiration and other biochemical processes, the third is the loss of ability of all cells to react to stimuli and the cessation cilia movements and heart beating.<sup>11,12</sup> In humans, minimal alveolar concentrations (MAC) concept was introduced

to characterize the potency of inhalational anesthetics.<sup>135</sup> At 0.1- 0.3 MAC, most humans show sensory distortions, memory loss and sleepiness; at 0.3- 0.5 MAC, responses to verbal commands cease and loss of consciousness occurs; at about 1.0 MAC, insensitivity to noxious stimuli occurs.<sup>68,135</sup> This marked concentration dependency of anesthetic potency suggests that some cells, or cellular processes, are more sensitive than others due to perhaps a higher number of the critical binding sites for the anesthetics. In neuronal assemblies, there are large numbers of excitable cells organized, via synaptic cell-cell adhesions, into higher order cellular assemblies. These communicate via action potentials and accomplish synchronous and coherent oscillations dependent on sensory stimuli. It is perhaps important that the plasma membrane at synapses is actively engaged in the endocytic vesicle recycling, amplifying the critical area of membranes significantly and underlying the high excitability of neuronal membranes.<sup>136-138</sup> The high electrical activity of neurons is related to their active maintenance of the physical properties of the lipid bilayer and the internal physico-chemical properties in the fluctuating cellular environments. Anesthetics and action potentials are highly relevant in this respect.<sup>8,136,139</sup> Surprisingly, although plant cells are sometimes considered not to be excitable,<sup>136,137</sup> the opposite is true.<sup>12,139-147</sup> In plant cells, action potentials, vesicle recycling, and sensitivity to anesthetics seem to be related to the active protection of the plasma membrane against ionic and structural disturbances.<sup>12,139,147-153</sup> These issues are prominent especially in neurons and specialized root apex cells both of which are very active in both electrical activities and endocytic vesicle recycling.<sup>143,147</sup> For both neurons and specialized plant cells, the endocytic (synaptic) recycling apparatus enhances the sensitivity to anesthetics due to large area of plasma membrane and the associated recycling vesicles, which is supported by the dynamic cytoskeleton.<sup>147,154-156</sup> It is emerging that these unique properties of brain neurons and plant root apex cells makes them exquisitely sensitive to anesthetics (Fig. 2).

Claude Bernard was the first to note that chloroplasts and their photosynthetic pathways are more sensitive to anesthetics than mitochondria and their respiratory pathways.<sup>12,25</sup> One of the striking differences between chloroplasts and mitochondria is stacking of internal chloroplast membranes into synaptic-like assemblies known as thylakoid grana which are essential for the photosynthetic pathways.<sup>157</sup> Intriguingly, in this respect, synaptic-like MORN-domain proteins have been reported for the thylakoid proteome.<sup>158</sup> On the other hand, internal membranes of mitochondria are much simpler and never form such prominent membrane



**Figure 2.** Both in animals and plants, organs with the highest activities of endocytic vesicle recycling and electric activities are implied in loss of consciousness (motility, sensitivity, and behavior).


stacks. Moreover, synaptic-like proteins including synaptotagmin-like E-SYT, yeast tricalbins and plant SYTs localize to membrane adhesions between intracellular organelles including plasma membranes, nuclei, mitochondria, chloroplasts, peroxisomes, ER membranes, and lipid bodies.<sup>159-161</sup> We are proposing that synaptic-like<sup>157</sup> stacking of thylakoid membranes into prominent chloroplast grana<sup>162,163</sup> is behind the higher sensitivity of the chloroplast photosynthesis to anesthetics<sup>25</sup> in comparison to the mitochondrial respiration (Fig. 3).

### Physical nature of anesthesia and consciousness

The unique aspect of anesthesia and consciousness is that these deep mysteries challenge both biology and physics. In fact, there are many aspects of the actions of anesthetics on all life which implicate a profound physical basis for both anesthesia and consciousness. Action potentials have not only electrical, but also mechanical aspects, as they change significantly plasma membrane thickness and even the length of electrically active neurons.<sup>164,167</sup> Moreover, protein activity is also highly dependent on physical factors. For example, the tertiary structures of proteins are easily deformed and this affects their activity.

Interestingly, action potentials also generate heat in excited membranes<sup>168-170</sup> which then unfolds proteins and fluidizes lipid bilayers.<sup>171</sup> Also anesthetics unfold proteins and induce fluidization of lipid bilayers.<sup>171-173</sup>





**Figure 3.** Claude Bernard discovered higher sensitivity of photosynthesis to anesthetics in comparison to respiration.<sup>12,25</sup> Chloroplast accomplish photosynthesis on stacked membranes know as thylakoid grana<sup>162,163</sup> which can be considered for inter-organelar synapses.<sup>157</sup> We propose that these stacked membranes are not only essential for photosynthesis but also makes this process more sensitive to anesthetics.

Importantly, the potency of anesthetics in inducing the loss of motility and consciousness decreases with the temperature increasing.<sup>174-176</sup>

Recent advances with xenon anesthesia strongly suggest that the currently favored proteins/ receptors model of anesthetics actions will require a serious update toward including the lipid bilayer as the primary target of xenon. This unique simple noble gas is chemically inert and acts on membranes only through its physical properties.<sup>177,178</sup> The physical nature of both anesthesia and consciousness is implicated also with the reversal of anesthetic induced loss of consciousness by high pressure. This pressure reversal was discovered by Johnson and Flagler who showed that anesthetized tadpoles regain activities at pressures of about 50 bar.<sup>178</sup> Since then this observation has been confirmed with many other organisms and also with diverse anesthetics. With respect to xenon, high pressure was shown to prevent free xenon diffusion within lipid bilayers and xenon was pushed out to accumulate in the middle of the lipid bilayer.<sup>177</sup> Furthermore, xenon was found to modulate the bilayer lateral pressure profile in a reversible fashion.<sup>178</sup> Xenon-induced changes to the critical proteins may be only the secondary consequences of these physical effects of xenon on lipid bilayers.<sup>178,180</sup> Alternatively, there might also be a genuine direct effect of xenon on proteins via their hydrophobic pockets. Intriguingly, xenon rapidly reverses electron spins and these electron spin effects were found to be different in *Drosophila* mutants which did not respond to the anesthetics.<sup>10</sup> The next perplexing finding is that both local and general

anesthetics show similar physical effects on membranes by lowering their melting temperatures and this anesthetic effect is reversed by high pressure.<sup>167,181,182</sup> There are further issues suggesting that physical phenomena are related to both anesthesia and consciousness based on quantum aspects of physical reality.<sup>7</sup> In conclusion, solving of these mysteries will be based on both, biology and physics, and relevant to both. This fundamental advance in our knowledge will help to understand the basic question of biology posed by the famous physicist Erwin Schrödinger: ‘what is life?’<sup>183</sup>

### The implications of a lack of a role for genes in consciousness

As noted above, the rapidity of the switching on and off of consciousness by anesthetics precludes a role for the necessarily slower action of gene expression. It can be argued that genes are in fact only necessary to 2 quite small, but nevertheless important, aspects of the life process, namely reproduction and the storage of necessary data (base sequence) to transcribe the peptides essential for the cell to function.<sup>184</sup> On this model, organisms as we know them today, were preceded by metabolizing proto-cells based on proteins; which regulated themselves in order to realize a proto-phenotype with the ability to engage in purposeful behavior. Such cells could not replicate themselves, but divided due to stresses on the cell membrane. The crucial step to true life as we know it was to recruit nucleobases to encode peptide sequences in DNA, thus allowing true replication and evolution to more complex organisms to commence. It has long been known that even the most primitive of organisms are capable of *purposeful* behavior<sup>185-187</sup> and that this can only be due to cellular proteins processing environmental information detected at the cell membrane.<sup>188</sup> Enucleated cells commutate with nucleated cells, obtaining small molecules to correct the deficient cells,<sup>189</sup> but are also able to survive up to several months<sup>190,191</sup> and organize their circadian clocks without any DNA and gene expression.<sup>133</sup> These considerations lend weight to the proposal that anesthetics act by disrupting the activity of proteins as enzymes,<sup>5</sup> specifically the (mostly not understood) processes in protein information processing, perhaps with those associated with membranes.

### Outlook

Although it is not clearly stated by most of the authors discussing consciousness, and some even claim that consciousness is an epiphenomenon, it is very obvious that consciousness is essential for survival and life in general.<sup>192,193</sup> Anyone of us losing consciousness would not

be able to survive a few days without the devoted assistance and help from our fellows. Similarly, life is not possible without recognizing danger via pain<sup>194</sup> and other kinds of negative experiences safeguarding survival,<sup>195-197</sup> all based on consciousness. It is obvious that consciousness is essential for any organism to have online access to their sensory information about their environments.<sup>193,198</sup> Importantly, consciousness gives all organisms ability to act as agents of their own interest<sup>199,200</sup> which is essential for their survival. This is essential for organisms to navigate successfully in complex environments that challenge their survival.

### Disclosure of potential conflicts of interest

No potential conflicts of interest were disclosed.

### References

- [1] Rudolph U, Antkowiak B. Molecular and neuronal substrates for general anaesthetics. *Nat Rev Neurosci* 2004; 5:709-72; <http://dx.doi.org/10.1038/nrn1496>
- [2] Perouansky M. The quest for a unified model of anesthetic action. A century in Claude Bernard's shadow. *Anesthesiology* 2012; 117:465-74; <http://dx.doi.org/10.1097/ALN.0b013e318264492e>
- [3] Rinaldi A. Reawakening anaesthesia research. *EMBO Rep* 2014; 15:1113-8; <http://dx.doi.org/10.15252/embr.201439593>
- [4] Bancroft WD, Richter GH. Claude Bernard's theory of narcosis. *Proc Natl Acad Sci USA* 1930; 16:573-577; <http://dx.doi.org/10.1073/pnas.16.9.573>
- [5] Franks NP, Lieb WR. Do general anaesthetics act by competitive binding to specific receptors? *Nature* 1984; 310:599-601; <http://dx.doi.org/10.1038/310599a0>
- [6] Hameroff SR. The entwined mysteries of anesthesia and consciousness: is there a common underlying mechanism? *Anesthesiology* 2006; 105:400-12; <http://dx.doi.org/10.1097/00000542-200608000-00024>
- [7] Hameroff SR, Craddock TJ, Tuszyński JA. Quantum effects in the understanding of consciousness. *J Integr Neurosci* 2014; 13:229-52; <http://dx.doi.org/10.1142/S0219635214400093>
- [8] Sonner JM. A hypothesis on the origin and evolution of the response to inhaled anesthetics. *Anesth Analg* 2008; 107:849-854; <http://dx.doi.org/10.1213/ane.0b013e31817ee684>
- [9] Sonner JM, Cantor RS. Molecular mechanisms of drug action: an emerging view. *Ann Rev Biophys* 2013; 42:143-67; <http://dx.doi.org/10.1146/annurev-biophys-083012-130341>
- [10] Turin L, Skoulakis EMC, Horsfield AP. Electron spin changes during general anesthesia in *Drosophila*. *Proc Natl Acad Sci USA* 2014; 111:E3524-33; <http://dx.doi.org/10.1073/pnas.1404387111>
- [11] Bernard C. Lectures on the Phenomena of Life Common to Animals and Plants. Charles C Thomas Pub Ltd (June 1974); 1878.
- [12] Grémiaux A, Yokawa K, Mancuso S, Baluška F. Plant anesthesia supports similarities between animals and plants: Claude Bernard's forgotten studies. *Plant Signal Behav* 2014; 9:e27886; <http://dx.doi.org/10.4161/psb.27886>
- [13] Lloyd FE. Some effects of narcotics on Spirogyra. *Anesth Analg* 1924; 3:9-19; <http://dx.doi.org/10.1213/00000539-192402000-00003>
- [14] White DC, Dundas CR. Effect of anaesthetics on emission of light by luminous bacteria. *Nature* 1970; 226:456-8; <http://dx.doi.org/10.1038/226456a0>
- [15] De Luccia TP. *Mimosa pudica*, *Dionaea muscipula* and anesthetics. *Plant Signal Behav* 2012; 7:1163-7; <http://dx.doi.org/10.4161/psb.21000>
- [16] Hall GM, Kirtland SJ, Baum H. The inhibition of mitochondrial respiration by inhalational anaesthetic agents. *Br J Anaesth* 1973; 45:1005-9; <http://dx.doi.org/10.1093/bja/45.10.1005>
- [17] Cohen PJ. Effect of anesthetics on mitochondrial function. *Anesthesiology* 1973; 39:153-64; <http://dx.doi.org/10.1097/00000542-197308000-00007>
- [18] Brunner EA, Cheng SC, Berman ML. Effects of anesthesia on intermediary metabolism. *Annu Rev Med* 1975; 26:391-401; <http://dx.doi.org/10.1146/annurev.me.26.020175.002135>
- [19] Nahrwold ML, Cohen PJ. Anesthetics and mitochondrial respiration. *Clin Anesth* 1975; 11:25-44
- [20] Nakao H, Ogi K, Yokono S, Ono J, Miyatake A. The effect of volatile anesthetics on light-induced phosphorylation in spinach chloroplasts. *Toxicol Lett* 1998; 100-101:135-8; [http://dx.doi.org/10.1016/S0378-4274\(98\)00177-5](http://dx.doi.org/10.1016/S0378-4274(98)00177-5)
- [21] La Monaca E, Fodale V. Effects of anesthetics on mitochondrial signaling and function. *Curr Drug Saf* 2012; 7:126-39; <http://dx.doi.org/10.2174/157488612802715681>
- [22] Ingram LO. Adaptation of membrane lipids to alcohols. *J Bacteriol* 1976; 125:670-8
- [23] Johnson SM, Saint John BE, Dine AP. Local anesthetics as antimicrobial agents: a review. *Surg Infect (Larchmt)* 2008; 9:205-13; <http://dx.doi.org/10.1089/sur.2007.036>
- [24] Weng Y, Yang L, Corringier PJ, Sonner JM. Anesthetic sensitivity of the *Gloeobacter violaceus* proton-gated ion channel. *Anesth Analg* 2010; 110:59-63; <http://dx.doi.org/10.1213/ANE.0b013e3181c4bc69>
- [25] Bernard C. Leçons sur les anesthésiques et sur l'asphyxie. Librairie J-B Baillière et Fils. 1875
- [26] Ewart AJ. On the Physics and Physiology of Protoplasmic Streaming in Plants. Oxford, 1903.
- [27] Osterhout WJV. Some aspects of protoplasmic motion. *J Gen Physiol* 1951; 35:519-27; <http://dx.doi.org/10.1085/jgp.35.3.519>
- [28] Garcia-Sierra F, Frixione E. Lidocaine, a local anesthetic, reversibly inhibits cytoplasmic streaming in *Vallisneria mesophyll* cells. *Protoplasma* 1993; 175:153-60; <http://dx.doi.org/10.1007/BF01385014>
- [29] Kiefer RT, Ploppa A, Krueger WA, Plank M, Nohé B, Haerberle HA, Unertl K, Dieterich HJ. Local anesthetics impair human granulocyte phagocytosis activity, oxidative burst, and CD11b expression in response to *Staphylococcus aureus*. *Anesthesiology* 2003; 98:842-8; <http://dx.doi.org/10.1097/00000542-200304000-00009>
- [30] Hameroff S. Why anesthetic mechanism research has failed, and what to do about it. <http://anesth.medicine.arizona.edu/system/files/pdfs>, Downloaded June 2015

- [31] Sonner JM. Issues in the design and interpretation of minimum alveolar anesthetic concentration (MAC) studies. *Anesth Analg* 2002; 95:609-14
- [32] Brosnan RJ, Yang L, Milutinovic PS, Zhao J, Laster MJ, Eger EI, 2nd, Sonner JM. Ammonia has anesthetic properties. *Anesth Analg* 2007; 104:430-3; <http://dx.doi.org/10.1213/01.ane.0000264072.97705.0f>
- [33] Yang L, Zhao J, Milutinovic PS, Brosnan RJ, Eger EI, 2nd, Sonner JM. Anesthetic properties of the ketone bodies  $\beta$ -hydroxybutyric acid and acetone. *Anesth Analg* 2007; 105:673-9; <http://dx.doi.org/10.1213/01.ane.0000278127.68312.dc>
- [34] Weng Y, Hsu TT, Zhao J, Nishimura S, Fuller GG, Sonner JM. Isovaleric, methylmalonic, and propionic acid decrease anesthetic EC50 in tadpoles, modulate glycine receptor function, and, interact with the lipid 1,2-dipalmitoyl-Sn-glycero-3-phosphocholine. *Anesth Analg* 2009; 108:1538-45; PMID:19372333; <http://dx.doi.org/10.1213/ane.0b013e31819cd964>
- [35] Eger EI, 2nd, Raines DE, Shafer SL, Hemmings HC, Jr, Sonner JM. Is a new paradigm needed to explain how inhaled anesthetics produce immobility? *Anesth Analg* 2008; 107:832-48; <http://dx.doi.org/10.1213/ane.0b013e318182aedb>
- [36] Tanaka K, Ludwig LM, Kersten JR, Pagel PS, Warltier DC. Mechanisms of cardioprotection, by volatile anesthetics. *Anesthesiology* 2004;100:707-21; PMID:15108989; <http://dx.doi.org/10.1097/00000542-200403000-00035>
- [37] De Hert SG, Turani F, Mathur S, Stowe DF. Cardioprotection with volatile anesthetics: mechanisms and clinical implications. *Anesth Analg* 2005; 100:1584-93; PMID:15920178; <http://dx.doi.org/10.1213/01.ANE.0000153483.61170.0C>
- [38] Agarwal B, Stowe DF, Dash RK, Bosnjak ZJ, Camara AK. Mitochondrial targets for volatile anesthetics against cardiac ischemia-reperfusion injury. *Front Physiol* 2014; 5:341; PMID:25278902; <http://dx.doi.org/10.3389/fphys.2014.00341>
- [39] Keller C, Grimm C, Wenzel A, Hafezi F, Remé C. Protective effect of halothane anesthesia on retinal light damage: inhibition of metabolic rhodopsin regeneration. *Invest Ophthalmol Vis Sci* 2001; 42:476-80
- [40] Stevenson GW, Hall S, Rudnick SJ, Alvord G, Rossio J, Urban W, Leventhal JB, Miller P, Seleny F, Stevenson HC. Halothane anesthesia decreases human monocyte hydrogen peroxide generation. Protection of monocytes by activation with gamma interferon. *Immunopharmacol Immunotoxicol* 1987; 9:489-510; PMID:3125239; <http://dx.doi.org/10.3109/08923978709035228>
- [41] Barodka VM, Acheampong E, Powell G, Lobach L, Logan DA, Parveen Z, Armstead V, Mukhtar M. Antimicrobial effects of liquid anesthetic isoflurane on *Candida albicans*. *J Transl Med* 2006; 4:46; PMID:17094810; <http://dx.doi.org/10.1186/1479-5876-4-46>
- [42] Manley SL. Phytochemistry of halomethanes: a production of selection or a metabolic accident? *Biogeochemistry* 2002; 60:163-80; <http://dx.doi.org/10.1023/A:1019859922489>
- [43] Laothawornkitkul J, Taylor JE, Paul ND, Hewitt CN. Biogenic volatile organic compounds in the Earth system. *New Phytol* 2009; 183:27-51; PMID:19422541; <http://dx.doi.org/10.1111/j.1469-8137.2009.02859.x>
- [44] Redeker KR, Wang N, Low JC, McMillan A, Tyler SC, Cicerone RJ. Emissions of methyl halides and methane from rice paddies. *Science* 2000; 290:966-9; PMID:11062125; <http://dx.doi.org/10.1126/science.290.5493.966>
- [45] Holopainen JK, Gershenzon J. Multiple stress factors and the emission of plant VOCs. *Trends Plant Sci* 2010; 15:176-84; PMID:20144557; <http://dx.doi.org/10.1016/j.tplants.2010.01.006>
- [46] Loreto F, Schnitzler JP. 2010. Abiotic stresses and induced BVOCs. *Trends Plant Sci* 15:154-66; PMID:20133178; <http://dx.doi.org/10.1016/j.tplants.2009.12.006>
- [47] Peñuelas J, Staudt M. BVOCs and global change. *Trends Plant Sci* 2010; 15:133-144; <http://dx.doi.org/10.1016/j.tplants.2009.12.005>
- [48] Kegge W, Pierik R. Biogenic volatile organic compounds and plant competition. *Trends Plant Sci* 2010; 15:126-32; PMID:20036599; <http://dx.doi.org/10.1016/j.tplants.2009.11.007>
- [49] Fang Z, Ionescu P, Chortkoff BS, Kandel L, Sonner J, Laster MJ, Eger EI, 2nd. Anesthetic potencies of n-alkanols: results of additivity and solubility studies suggest a mechanism of action similar to that for conventional inhaled anesthetics. *Anesth Analg* 1997; 84:1042-8; PMID:9141929; <http://dx.doi.org/10.1213/00000539-199705000-00017>
- [50] Hau KM, Connell DW, Richardson BJ. A study of the biological partitioning behavior of n-alkanes and n-alkanols in causing anesthetic effects. *Regul Toxicol Pharmacol* 2002; 35:273-9; <http://dx.doi.org/10.1006/rtp.2001.1531>
- [51] Butterbach-Bahl K, Baggs EM, Dannenmann M, Kiese R, Zechmeister-Boltenstern S. Nitrous oxide emissions from soils: how well do we understand the processes and their controls? *Philos Trans R Soc Lond B Biol Sci* 2013; 368:20130122; PMID:23713120; <http://dx.doi.org/10.1098/rstb.2013.0122>
- [52] Könneke M, Schubert DM, Brown PC, Hügler M, Standfest S, Schwander T, Schada von Borzyskowski L, Erb TJ, Stahl DA, Berg IA. Ammonia-oxidizing archaea use the most energy-efficient aerobic pathway for CO<sub>2</sub> fixation. *Proc Natl Acad Sci USA* 2014; 111:8239-44; PMID:24843170; <http://dx.doi.org/10.1073/pnas.1402028111>
- [53] Hu HW, Chen D, He JZ. Microbial regulation of terrestrial nitrous oxide formation: understanding the biological pathways for prediction of emission rates. *FEMS Microbiol Rev* 2015; 39:729-49
- [54] Howe GA, Schillmiller AL. Oxylipin metabolism in response to stress. *Curr Opin Plant Biol* 2002; 5:230-6; PMID:11960741; [http://dx.doi.org/10.1016/S1369-5266\(02\)00250-9](http://dx.doi.org/10.1016/S1369-5266(02)00250-9)
- [55] Chechetkin IR, Mukhitova FK, Blufard AS, Yarin AY, Antsygina LL, Grechkin AN. Unprecedented pathogen-inducible complex oxylipins from flax linolipins A and B. *FEBS J* 2009; 276:4463-72; PMID:19645727; <http://dx.doi.org/10.1111/j.1742-4658.2009.07153.x>
- [56] Wang F, Cui X, Sun Y, Dong CH. Ethylene signaling and regulation in plant growth and stress responses. *Plant Cell Rep* 2013; 32:1099-109; PMID:23525746; <http://dx.doi.org/10.1007/s00299-013-1421-6>



- [57] Kazan K. Diverse roles of jasmonates and ethylene in abiotic stress tolerance. *Trends Plant Sci* 2015; 20:219-29; PMID:25731753; <http://dx.doi.org/10.1016/j.tplants.2015.02.001>
- [58] Calixto JB, Beirith A, Ferreira J, Santos AR, Filho VC, Yunes RA. Naturally occurring antinociceptive substances from plants. *Phytother Res* 2000; 14:401-18; PMID:10960893; [http://dx.doi.org/10.1002/1099-1573\(200009\)14:6%3c401::AID-PTR762%3e3.0.CO;2-H](http://dx.doi.org/10.1002/1099-1573(200009)14:6%3c401::AID-PTR762%3e3.0.CO;2-H)
- [59] Askitopoulou H, Ramoutsas IA, Konsolaki E. Analgesia and anesthesia: etymology and literary history of related greek words. *Anesth Analg* 2000; 91:486-91; PMID:10910873
- [60] Kennedy DO, Wightman EL. Herbal extracts and phytochemicals: plant secondary metabolites and the enhancement of human brain function. *Adv Nutr* 2011; 2:32-50; PMID:22211188; <http://dx.doi.org/10.3945/an.110.000117>
- [61] Kennedy DO. *Plants and the Human Brain*. Oxford University Press, 2014
- [62] Meotti FC, Lemos de Andrade E, Calixto JB. TRP modulation by natural compounds. *Handb Exp Pharmacol* 2014; 223:1177-238; PMID:24961985; [http://dx.doi.org/10.1007/978-3-319-05161-1\\_19](http://dx.doi.org/10.1007/978-3-319-05161-1_19)
- [63] Dolferus R, Jacobs M, Peacock WJ, Dennis ES. Differential interactions of promoter elements in stress responses of the Arabidopsis Adh gene. *Plant Physiol* 1994; 105:1075-87; PMID:7972489; <http://dx.doi.org/10.1104/pp.105.4.1075>
- [64] Strommer J. The plant ADH gene family. *Plant J* 2011; 66:128-42; PMID:21443628; <http://dx.doi.org/10.1111/j.1365-313X.2010.04458.x>
- [65] Taylorson RB, Hendricks SB. Overcoming dormancy, in seeds with ethanol and other anesthetics. *Planta* 1979; 145:507-10; PMID:24317868
- [66] Dillard MM. Ethylene – the new general anesthetic. *J Natl Med Assoc* 1930; 22:10-11; PMID:20892343
- [67] Campagna JA, Miller KW, Forman SA. Mechanisms of actions of inhaled anesthetics. *New Engl J Med* 2003; 348:2110-24; PMID:12761368; <http://dx.doi.org/10.1056/NEJMra021261>
- [68] Kopp Lugli A, Yost CS, Kindler CH. Anaesthetic mechanisms: update on the challenge of unravelling the mystery of anaesthesia. *Eur J Anaesthesiol* 2009; 26:807-20; PMID:19494779; <http://dx.doi.org/10.1097/EJA.0b013e32832d6b0f>
- [69] Freebairn HT, Buddendenhagen IW. Ethylene production by *Pseudomonas solanacearum*. *Nature* 1964; 202:313-4; PMID:14167811; <http://dx.doi.org/10.1038/202313a0>
- [70] Smith AM. Ethylene as a cause of soil fungistasis. *Nature* 1973; 246:311-3; PMID:4586317; <http://dx.doi.org/10.1038/246311a0>
- [71] Lynch JM, Harper SHT. Formation of ethylene by a soil fungus. *J Gen Microbiol* 1974; 80:187-95; <http://dx.doi.org/10.1099/00221287-80-1-187>
- [72] Smith AM, James Cook R. Implications of ethylene production by bacteria for biological balance of soil. *Nature* 1974; 252:703-5; PMID:4437621; <http://dx.doi.org/10.1038/252703b0>
- [73] Primrose SB, Dilworth MJ. Ethylene production by bacteria. *J Gen Microbiol* 1976; 93:177-81; PMID:772166; <http://dx.doi.org/10.1099/00221287-93-1-177>
- [74] Considine PJ, Flynn N, Patching JW. Ethylene production by soil microorganisms. *Appl Environ Microbiol* 1977; 33:977-9; PMID:869541
- [75] Graham JH, Linderman RG. Ethylene production by ectomycorrhizal fungi, *Fusarium oxysporum* f. sp. pini, and by aseptically synthesized ectomycorrhizae and Fusarium-infected Douglas-fir roots. *Can J Microbiol* 1980; 26:1340-7; PMID:7214223; <http://dx.doi.org/10.1139/m80-222>
- [76] Huang TC, Chow TJ. Ethylene production by blue-green algae. *Bot Bull Acad Sinica* 1984; 25:81-6
- [77] Kauppi N, Kauppi A, Garty J. Ethylene produced by the lichen *Cladonia stellaris* exposed to sulphur and heavy metal-containing solutions under acidic conditions. *New Phytol* 1998; 139:537-47; <http://dx.doi.org/10.1046/j.1469-8137.1998.00214.x>
- [78] Chagué V, Elad Y, Barakat R, Tudzynski P, Sharon A. Ethylene biosynthesis in *Botrytis cinerea*. *FEMS Microbiol Ecol* 2002; 40:143-9; <http://dx.doi.org/10.1111/j.1574-6941.2002.tb00946.x>
- [79] Plettner I, Steinke M, Malin G. Ethene (ethylene) production in the marine macroalga *Ulva (Enteromorpha) intestinalis* L. (Chlorophyta, Ulvophyceae): effect of light-stress and co-production with dimethyl sulphide. *Plant Cell Environ* 2005; 28:1136-45; <http://dx.doi.org/10.1111/j.1365-3040.2005.01351.x>
- [80] Quadir A, Hewett EW, Long PG, Dilley DR. A non-ACC pathway for ethylene biosynthesis in *Botrytis cinerea*. *Postharv Biol Technol* 2011; 62:314-8; <http://dx.doi.org/10.1016/j.postharvbio.2011.06.003>
- [81] Whalen FX, Bacon DR, Smith HM. Inhaled anesthetics: an historical overview. *Best Pract Res Clin Anaesthesiol* 2005; 19:323-30; PMID:16013684; <http://dx.doi.org/10.1016/j.bpa.2005.02.001>
- [82] Mazurek MJ. Dr. Chauncey Leake and the development of divinyl oxide from bench to bedside. *CSA Bull* 2007; 86-9
- [83] Finer B. Divinyl ether. *Brit J Anaesth* 1965; 37:661-6; PMID:5320087; <http://dx.doi.org/10.1093/bja/37.9.661>
- [84] Itoh A, Howe GA. Molecular cloning of a divinyl ether synthase. *J Biol Chem* 2001; 276:3620-27; PMID:11060314; <http://dx.doi.org/10.1074/jbc.M008964200>
- [85] Stumpe M, Carsjens JG, Göbel C, Feussner I. Divinyl ether synthesis in garlic bulbs. *J Exp Bot* 2008; 59:907-15; PMID:18326559; <http://dx.doi.org/10.1093/jxb/ern010>
- [86] Fammartino A, Verdaguer B, Fournier J, Tamietti G, Carbonne F, Esquerré-Tugayé MT, Cardinale F. Coordinated transcriptional regulation of the divinyl ether biosynthetic genes in tobacco by signal molecules related to defense. *Plant Physiol Biochem* 2010; 48:225-31; PMID:20137961; <http://dx.doi.org/10.1016/j.plaphy.2010.01.012>
- [87] Fammartino A, Cardinale F, Göbel C, Mène-, Saffrané L, Fournier J, Feussner I, Esquerré-Tugayé MT. Characterization of a divinyl ether biosynthetic pathway specifically associated with pathogenesis in tobacco. *Plant Physiol* 2007; 143:378-88; PMID:17085514; <http://dx.doi.org/10.1104/pp.106.087304>
- [88] Dasta J. Local anesthetics: evolving to a new standard of care. *Pharm Pract New Spec Rep* 2013; January:1-8.
- [89] Golembiewski J, Dasta J. Evolving role of local anesthetics in managing postsurgical analgesia. *Clin*

- Therap 2015; 37:1354-1371; <http://dx.doi.org/10.1016/j.clinthera.2015.03.017>
- [90] Behçet Al. The source-synthesis: history and use of atropine. *J Acad Emerg Med* 2014; 13:2-3; <http://dx.doi.org/10.5152/jaem.2014.1120141>
- [91] Façanha MF, Gomes LC. Efficacy of menthol as an anesthetic for tambaqui (*Colossoma macropomum*, Characiformes: Characidae). *Acta Amazon* 2005; 35:71-5; <http://dx.doi.org/10.1590/S0044-59672005000100011>
- [92] Watt EE, Betts BA, Kotey FO, Humbert DJ, Griffith TN, Kelly EW, Veneskey KC, Gill N, Rowan KC, Jenkins A, Hall AC. Menthol shares general anesthetic activity and sites of action on the GABA(A) receptor with the intravenous agent, propofol. *Eur J Pharmacol* 2008; 590:120-6; PMID:18593637; <http://dx.doi.org/10.1016/j.ejphar.2008.06.003>
- [93] Ghelardini C, Galeotti N, Mazzanti G. Local anaesthetic activity of monoterpenes and phenylpropanes of essential oils. *Planta Med* 2001; 67:564-6; PMID:11509984; <http://dx.doi.org/10.1055/s-2001-16475>
- [94] de Lima Silva L, Thomas da Silva D, Garlet QI, Cunha MA, Mallmann CA, Baldisserotto B, Longhi SJ, Soares Pereira AM, Heinzmann BA. Anesthetic activity of Brazilian native plants in silver catfish (*Rhamdia quelen*). *Neotrop Ichthyol* 2013; 11:443-51; <http://dx.doi.org/10.1590/S1679-62252013000200014>
- [95] Burton RR. G-induced loss of consciousness: definition, history, current status. *Aviat Space Environ Med* 1988; 59:2-5; PMID:3281645
- [96] Johanson DC, Pheeny HT. A new look at the loss of consciousness experience within the US. *Naval forces. Aviat Space Environ Med* 1988; 59:6-8; PMID:3355469
- [97] Diehl RR. Vasovagal syncope and Darwinian fitness. *Clin Auton Res* 2005; 15:126-129; PMID:15834770; <http://dx.doi.org/10.1007/s10286-005-0244-0>
- [98] Alboni P, Alboni M. Vasovagal syncope as a manifestation of an evolutionary selected trait. *J Atrial Fibril* 2014; 7:97-102
- [99] Rose da Silva MFL. Syncope: epidemiology, etiology, and prognosis. *Front Physiol* 2014; 5:4; PMID:24478717
- [100] Palmer LK, Rannels SL, Kimball SR, Jefferson LS, Keil RL. Inhibition of mammalian translation initiation by volatile anesthetics. *Am J Physiol Endocrinol Metab* 2006; 290:E1267-75; PMID:16434554; <http://dx.doi.org/10.1152/ajpendo.00463.2005>
- [101] Uesono Y. Environmental stresses and clinical drugs paralyze a cell. *Commun Integr Biol* 2009; 2:275-8; PMID:19641750; <http://dx.doi.org/10.4161/cib.2.3.8226>
- [102] Uesono Y, Toh EA, Kikuchi Y, Terashima I. Structural analysis of compounds with actions similar to local anesthetics and antipsychotic phenothiazines in yeast. *Yeast* 2011; 28:391-404; PMID:21374719; <http://dx.doi.org/10.1002/yea.1846>
- [103] Safford WE. An Aztec narcotic. *J Hered* 1915; 6:291-311
- [104] Osmond H. Ololiuqui: The ancient Aztec narcotic. *J Men Dis* 1955; 101:526-37
- [105] Spiller HA, Hale JR, De Boer JZ. The Delphic Oracle: A multidisciplinary defense of the gaseous vent theory. *Clin Toxicol* 2002; 40:189-96
- [106] Icaza EE, Mashour GA. Altered states. *Psychedelics and anesthetics. Anesthesiology* 2013; 119:1255-60; PMID:24061599; <http://dx.doi.org/10.1097/01.anes.000435635.42332.ee>
- [107] Dudley R. Ethanol, fruit ripening, and the historical origins of human alcoholism in primate frugivory. *Integr Comp Biol* 2004; 44:315-23; PMID:21676715; <http://dx.doi.org/10.1093/icb/44.4.315>
- [108] McGovern PE, Mirzoian A, Hall GR. Ancient Egyptian herbal wines. *Proc Natl Acad Sci USA* 2013; 106:7361-6; <http://dx.doi.org/10.1073/pnas.0811578106>
- [109] Carrigan MA, Uryasev O, Frye CB, Eckman BL, Myers CR, Hurley TD, Benner SA. Hominids adapted to metabolize ethanol long before human-directed fermentation. *Proc Natl Acad Sci USA* 2015; 112:458-63; PMID:25453080; <http://dx.doi.org/10.1073/pnas.1404167111>
- [110] Hockings KJ, Bryson-Morrison N, Carvalho S, Fujisawa M, Humle T, McGrew WC, Nakamura M, Ohashi G, Yamanashi Y, Yamakoshi G, et al. Tools to tipple: ethanol ingestion by wild chimpanzees using leaf-sponges. *R Soc Open Sci* 2015; 2:150150; PMID:26543588; <http://dx.doi.org/10.1098/rsos.150150>
- [111] Wiens F, Zitzmann A, Lachance MA, Yegles M, Pragst F, Wurst FM, von Holst D, Guan SL, Spanagel R. Chronic intake of fermented floral nectar by wild treeshrews. *Proc Natl Acad Sci USA* 2008; 105:10426-31; PMID:18663222; <http://dx.doi.org/10.1073/pnas.0801628105>
- [112] Devineni AV, Heberlein U. Addiction-like behavior in *Drosophila*. *Commun Integr Biol* 2010; 3:357-9; PMID:20798826; <http://dx.doi.org/10.4161/cib.3.4.11885>
- [113] Shohat-Ophir G, Kaun KR, Azanchi R, Mohammed H, Heberlein U. Sexual deprivation increases ethanol intake in *Drosophila*. *Science* 2012; 335:1351-5; PMID:22422983; <http://dx.doi.org/10.1126/science.1215932>
- [114] Peru Y, Colón de Portugal RL, Ojelade SA, Penninti PS, Dove RJ, Nye MJ, Acevedo SF, Lopez A, Rodan AR, Rothenfluh A. Long-lasting, experience-dependent alcohol preference in *Drosophila*. *Addict Biol* 2014; 19:392-401; PMID:24164972; <http://dx.doi.org/10.1111/adb.12105>
- [115] Lydic R, Baghdoyan HA. Sleep, anesthesiology, and the neurobiology of arousal state control. *Anesthesiology* 2005; 103:1268-95; PMID:16306742; <http://dx.doi.org/10.1097/00000542-200512000-00024>
- [116] Alkire MT, Hudetz AG, Tononi G. Consciousness and anesthesia. *Science* 2008; 322:876-80; PMID:18988836; <http://dx.doi.org/10.1126/science.1149213>
- [117] Hutt A. Sleep and anesthesia. *Front Neurosci* 2009; 3:408-9
- [118] Franks NP. General anaesthesia: from molecular targets to neuronal pathways of sleep and arousal. *Nat Rev Neurosci* 2008; 9:370-86; PMID:18425091; <http://dx.doi.org/10.1038/nrn2372>
- [119] Franks NP, Zecharia AY. Sleep and general anesthesia. *Can J Anaesth* 2011; 58:139-48; PMID:21170623; <http://dx.doi.org/10.1007/s12630-010-9420-3>
- [120] McDermott CM, LaHoste GJ, Chen C, Musto A, Bazan NG, Magee JC. Sleep deprivation causes behavioral, synaptic, and membrane excitability alterations in hippocampal neurons. *J Neurosci* 2003; 23:9687-95; PMID:14573548

- [121] Walker MP. The role of sleep in cognition and emotion. *Ann NY Acad Sci* 2009; 1156:168-97; PMID:19338508; <http://dx.doi.org/10.1111/j.1749-6632.2009.04416.x>
- [122] Rasch B, Born J. About sleep's role in memory. *Physiol Rev* 2013; 93:681-766; PMID:23589831; <http://dx.doi.org/10.1152/physrev.00032.2012>
- [123] Hobson JA. REM sleep and dreaming: towards a theory of protoconsciousness. *Nat Rev Neurosci* 2009; 10:803-13; PMID:19794431
- [124] Choi S, Yu E, Lee S, Llinás RR. Altered thalamocortical rhythmicity and connectivity in mice lacking CaV3.1 T-type Ca<sup>2+</sup> channels in unconsciousness. *Proc Natl Acad Sci USA* 2015; pii: 201420983
- [125] Kottler B, Bao H, Zalucki O, Imlach W, Troup M, van Alphen B, Paulk A, Zhang B, van Swinderen B. A sleep/wake circuit controls isoflurane sensitivity in *Drosophila*. *Curr Biol* 2013; 23:594-8; PMID:23499534; <http://dx.doi.org/>; <http://dx.doi.org/10.1016/j.cub.2013.02.021>
- [126] Price DD. Psychological and neural mechanisms of the affective dimension of pain. *Science* 2000; 288:1769-72; PMID:10846154; <http://dx.doi.org/10.1126/science.288.5472.1769>
- [127] Wager TD, Atlas LY. How Is Pain influenced by cognition? Neuroimaging weighs in. *Persp Psych Sci* 2013; 8:91-7; <http://dx.doi.org/10.1177/1745691612469631>
- [128] Zeidan F, Martucci KT, Kraft RA, McHaffie JG, Coghill RC. Neural correlates of mindfulness meditation-related anxiety relief. *Soc Cogn Affect Neurosci* 2014; 9:751-9; PMID:23615765; <http://dx.doi.org/10.1093/scan/nst041>
- [129] Jensen KB, Kaptchuk TJ, Kirsch I, Raicek J, Lindstrom KM, Berna C, Gollub RL, Ingvar M, Kong J. Nonconscious activation of placebo and nocebo pain responses. *Proc Natl Acad Sci USA* 2012; 109:15959-64; PMID:23019380; <http://dx.doi.org/10.1073/pnas.1202056109>
- [130] Jensen K, Kirsch I, Odmalm S, Kaptchuk TJ, Ingvar M. Classical conditioning of analgesic and hyperalgesic pain responses without conscious awareness. *Proc Natl Acad Sci USA* 2015; 112:7863-7; <http://dx.doi.org/10.1073/pnas.1504567112>
- [131] Price DD. Unconscious and conscious mediation of analgesia and hyperalgesia. *Proc Natl Acad Sci USA* 2015; 112:7624-5; PMID:26056258; <http://dx.doi.org/10.1073/pnas.1508765112>
- [132] Brown EN, Lydic R, Schiff ND. General anesthesia, sleep, and coma. *N Engl J Med* 2010; 363:2638-50; PMID:21190458; <http://dx.doi.org/10.1056/NEJMra0808281>
- [133] O'Neill JS, Reddy AB. Circadian clocks in human red blood cells. *Nature* 2011; 469:498-503; PMID:21270888; <http://dx.doi.org/10.1038/nature09702>
- [134] Nakajima M, Imai K, Ito H, Nishiwaki T, Murayama Y, Iwasaki H, Oyama T, Kondo T. Reconstitution of circadian oscillation of cyanobacterial KaiC phosphorylation in vitro. *Science* 2005; 308:414-15; PMID:15831759; <http://dx.doi.org/10.1126/science.1108451>
- [135] Eger EI, Saidman LJ, Brandstater B. Minimum alveolar anesthetic concentration: a standard of anesthetic potency. *Anesthesiology* 1965; 26:756-63; PMID:5844267; <http://dx.doi.org/10.1097/00000542-196511000-00010>
- [136] Cook ND. The neuron-level phenomena underlying cognition and consciousness: synaptic activity and the action potential. *Neuroscience* 2008; 153:556-70; PMID:18406536; <http://dx.doi.org/10.1016/j.neuroscience.2008.02.042>
- [137] Cook ND, Carvalho GB, Damasio A. From membrane excitability to metazoan psychology. *Trends Neurosci* 2014; 37:698-705; PMID:25176475; <http://dx.doi.org/10.1016/j.tins.2014.07.011>
- [138] Llinás RR. Intrinsic electrical properties of mammalian neurons and CNS function: a historical perspective. *Front Cell Neurosci* 2014; 8:320
- [139] Baluška F, Wan YL. Physical control over endocytosis. In: *Endocytosis in Plants*. Šamaj J. (ed), Springer Verlag; 2012.
- [140] Wayne R. Excitability in plant cells. *Am Sci* 1993; 81:140-51.
- [141] Wayne R. The excitability of plant cells: With a special emphasis on Characean internodal cells. *Bot Rev* 1994; 60:265-7; PMID:11539934; <http://dx.doi.org/10.1007/BF02960261>
- [142] Beilby MJ. Action potential in Charophytes. *Int Rev Cytol* 2007; 257:43-83; PMID:17280895; [http://dx.doi.org/10.1016/S0074-7696\(07\)57002-6](http://dx.doi.org/10.1016/S0074-7696(07)57002-6)
- [143] Masi E, Ciszak M, Stefano G, Renna L, Azzarello E, Pandolfi C, Mugnai S, Baluška F, Arecchi FT, Mancuso S. Spatio-temporal dynamics of the electrical network activity in the root apex. *Proc Natl Acad Sci USA* 2009; 106:4048-53; PMID:19234119; <http://dx.doi.org/10.1073/pnas.0804640106>
- [144] Hedrich R. Ion channels in plants. *Physiol Rev* 2012; 92:1777-811; PMID:23073631; <http://dx.doi.org/10.1152/physrev.00038.2011>
- [145] Król E, Dziubinska H, Trębacz K. What do plants need action potentials for? In: *Action Potential: Biophysical and Cellular Context, Initiation, Phases and Propagation*. DuBois ML. (ed), 2010; 1-26
- [146] Volkov A. *Plant Electrophysiology: Signaling and Responses*. Springer Verlag; 2012.
- [147] Baluška F, Mancuso S. Root apex transition zone as oscillatory zone. *Front Plant Sci* 2013; 4:354
- [148] Goldsworthy A. The evolution of plant action potentials. *J Theor Biol* 1983; 103:645-8; [http://dx.doi.org/10.1016/0022-5193\(83\)90287-4](http://dx.doi.org/10.1016/0022-5193(83)90287-4)
- [149] Steinhardt RA, Bi G, Alderton JM. Cell membrane resealing by a vesicular, mechanism similar to neurotransmitter release. *Science* 1994; 263:390-3; PMID:7904084; <http://dx.doi.org/10.1126/science.7904084>
- [150] Andrews NW, Chakrabarti S. There's more to life than neurotransmission: the regulation of exocytosis by synaptotagmin VII. *Trends Cell Biol* 2005; 15:626-31; PMID:16168654; <http://dx.doi.org/10.1016/j.tcb.2005.09.001>
- [151] Fu D, Vissavajhala P, Hemmings HC Jr. Volatile anaesthetic effects on phospholipid binding to synaptotagmin 1, a presynaptic Ca<sup>2+</sup> sensor. *Br J Anaesth* 2005; 95:216-21; PMID:15923266; <http://dx.doi.org/10.1093/bja/aei163>
- [152] Schapire AL, Valpuesta V, Botella MA. Plasma membrane repair in plants. *Trends Plant Sci* 2009; 14:645-52; PMID:19819752; <http://dx.doi.org/10.1016/j.tplants.2009.09.004>
- [153] Schapire AL, Voigt B, Jasik J, Rosado A, Lopez-Cobollo R, Menzel D, Salinas J, Mancuso S, Valpuesta V, Baluška



- F, Botella MA. *Arabidopsis* synaptotagmin 1 is required for the maintenance of plasma membrane integrity and cell viability. *Plant Cell* 2009; 20:3374-88; <http://dx.doi.org/10.1105/tpc.108.063859>
- [154] Morrow IC, Parton RG. Flotillins and the PHB domain protein family: rafts, worms and anaesthetics. *Traffic* 2005; 6:725-40; PMID:16101677; <http://dx.doi.org/10.1111/j.1600-0854.2005.00318.x>
- [155] Pristerá A, Okuse K. Building excitable membranes: lipid rafts and multiple controls on trafficking of electrogenic molecules. *Neuroscientist* 2012; 18:70-81; <http://dx.doi.org/10.1177/1073858410393977>
- [156] Zhao X, Li R, Lu C, Baluška F, Wan Y. Di-4-ANEPPDHQ, a fluorescent probe for the visualisation of membrane microdomains in living *Arabidopsis thaliana* cells. *Plant Physiol Biochem* 2015; 87:53-60; PMID:25549979; <http://dx.doi.org/10.1016/j.plaphy.2014.12.015>
- [157] Baluška F, Mancuso S. Synaptic view of eukaryotic cell. *Int J Gen Syst* 2014; 43:740-56; <http://dx.doi.org/10.1080/03081079.2014.920999>
- [158] Peltier JB, Ytterberg AJ, Sun Q, van Wijk KJ. New functions of the thylakoid membrane proteome of *Arabidopsis thaliana* revealed by a simple, fast, and versatile fractionation strategy. *J Biol Chem* 2004; 279:49367-83; PMID:15322131; <http://dx.doi.org/10.1074/jbc.M406763200>
- [159] Toulmay A, Prinz WA. A conserved membrane-binding domain targets proteins to organelle contact sites. *J Cell Sci* 2012; 125:49-58; PMID:22250200; <http://dx.doi.org/10.1242/jcs.085118>
- [160] Pérez-Lara A, Jahn R. Extended synaptotagmins (E-Syts): architecture and dynamics of membrane contact sites revealed. *Proc Natl Acad Sci USA* 2015; 112:4837-8; <http://dx.doi.org/10.1073/pnas.1504487112>
- [161] Pérez-Sancho J, Vanneste S, Lee E, McFarlane HE, Esteban Del Valle A, Valpuesta V, Friml J, Botella MA, Rosado A. The *Arabidopsis* Synaptotagmin1 is enriched in endoplasmic reticulum-plasma membrane contact sites and confers cellular resistance to mechanical stresses. *Plant Physiol* 2015; 168:132-43; <http://dx.doi.org/10.1104/pp.15.00260>
- [162] Mustardy L, Garab G. Granum revisited. A three-dimensional model – where things fall into place. *Trends Plant Sci* 2003; 8: 117-22; PMID:12663221; [http://dx.doi.org/10.1016/S1360-1385\(03\)00015-3](http://dx.doi.org/10.1016/S1360-1385(03)00015-3)
- [163] Kim EH, Chow WS, Horton P, Anderson JM. Entropy-assisted stacking of thylakoid membranes. *Biochim Biophys Acta* 2005; 1708:187-95; PMID:15953475; <http://dx.doi.org/10.1016/j.bbabi.2005.03.011>
- [164] Iwasa K, Tasaki I. Mechanical changes in squid giant axons associated with production of action potentials. *Biochem Biophys Res Commun* 1980; 95:1328-31; [http://dx.doi.org/10.1016/0006-291X\(80\)91619-8](http://dx.doi.org/10.1016/0006-291X(80)91619-8)
- [165] Iwasa K, Tasaki I, Gibbons RC. Swelling of nerve fibres associated with action potentials. *Science* 1980; 210:338-9; PMID:7423196; <http://dx.doi.org/10.1126/science.7423196>
- [166] Terakawa S. Changes in intracellular pressure in squid giant axons associated with production of action potentials. *Biochem Biophys Res Commun* 1983; 114:1006-10; PMID:6615499; [http://dx.doi.org/10.1016/0006-291X\(83\)90661-7](http://dx.doi.org/10.1016/0006-291X(83)90661-7)
- [167] Heimburg T. Nerves and anaesthesia: A physics perspective on medicine. *Anesth Analg* 2014; 8:252-63
- [168] Abbott BC, Hill AV, Howarth JV. The positive and negative heat production associated with a nerve impulse. *Proc Roy Soc London B* 1958; 148:149-87; <http://dx.doi.org/10.1098/rspb.1958.0012>
- [169] Ritchie JM, Keynes RD. The production and absorption of heat associated with electrical activity in nerve and electric organ. *Q Rev Biophys* 1985; 18:451-76; PMID:3916342; <http://dx.doi.org/10.1017/S0033583500005382>
- [170] Tasaki I, Byrne PM. Heat production associated with synaptic transmission in the bullfrog spinal cord. *Brain Res* 1987; 407:386-9; PMID:3032367; [http://dx.doi.org/10.1016/0006-8993\(87\)91119-X](http://dx.doi.org/10.1016/0006-8993(87)91119-X)
- [171] Ueda I, Yoshida T. Hydration of lipid membranes and the action mechanisms of anesthetics and alcohols. *Chem Phys Lip* 1999; 101:65-79; [http://dx.doi.org/10.1016/S0009-3084\(99\)00056-0](http://dx.doi.org/10.1016/S0009-3084(99)00056-0)
- [172] Papahadjopoulos D, Jacobson K, Poste G, Shepherd G. Effects of local anesthetics on membrane properties. I. Changes in the fluidity of phospholipid bilayers. *Biochim Biophys Acta* 1975; 394: 504-19; PMID:1148230; [http://dx.doi.org/10.1016/0005-2736\(75\)90137-6](http://dx.doi.org/10.1016/0005-2736(75)90137-6)
- [173] Zapata-Morin PA, Sierra-Valdez FJ, Ruiz-Suárez JC. The interaction of local anesthetics with lipid membranes. *J Mol Graph Model* 2014; 53:200-5; PMID:25181454; <http://dx.doi.org/10.1016/j.jmkgm.2014.08.001>
- [174] Cherkin A, Catchpool JF. Temperature dependence of anesthesia in goldfish. *Science* 1964; 144:1460-2; PMID:14171540; <http://dx.doi.org/10.1126/science.144.3625.1460>
- [175] Martin BJ. Evaluation of hypothermia for anesthesia in reptiles and amphibians. *ILAR J* 1995; 37:186-90; PMID:11528038; <http://dx.doi.org/10.1093/ilar.37.4.186>
- [176] Franks NP, Lieb WR. Temperature dependence of the potency of volatile general anesthetics: implications for in vitro experiments. *Anesthesiology* 1996; 84:716-20; PMID:8659800; <http://dx.doi.org/10.1097/0000542-199603000-00027>
- [177] Yamamoto E, Akimoto T, Shimizu H, Hirano Y, Yasui M, Yasuoka K. Diffusive nature of xenon anesthetic changes properties of a lipid bilayer: molecular dynamics simulations. *J Phys Chem B* 2012; 116:8989-95; PMID:22715916; <http://dx.doi.org/10.1021/jp303330c>
- [178] Booker RD, Sum AK. Biophysical changes induced by xenon on phospholipid bilayers. *Biochim Biophys Acta* 2013; 1828:1347-56; PMID:23376329; <http://dx.doi.org/10.1016/j.bbame.2013.01.016>
- [179] Johnson FH, Flagler EA. Hydrostatic pressure reversal of narcosis in tadpoles. *Science* 1950; 112:91-2; PMID:15442245; <http://dx.doi.org/10.1126/science.112.2899.91-a>
- [180] Lundbaek JA, Collingwood SA, Ingólfsson HI, Kapoor R, Andersen OS. Lipid bilayer regulation of membrane protein function: gramicidin channels as molecular force probes. *J R Soc Interface* 2010; 7:373-95; PMID:19940001; <http://dx.doi.org/10.1098/rsif.2009.0443>
- [181] Heimburg T, Jackson AD. The thermodynamics of general anesthesia. *Biophys J* 2007; 92:3159-65; PMID:17293400; <http://dx.doi.org/10.1529/biophysj.106.099754>
- [182] Graesbøll K, Sasse-Middelhoff H, Heimburg T. The thermodynamics of general and local anesthesia.



- Biophys J 2014; 106:2143-56; PMID:24853743; <http://dx.doi.org/10.1016/j.bpj.2014.04.014>
- [183] Schrödinger E. What Is Life? The Physical Aspect of the Living Cell. Based on lectures delivered under the auspices of the Dublin Institute for Advanced Studies at Trinity College, Dublin, in February 1943. Cambridge University Press; 1944.
- [184] Annala A, Baverstock K. Genes without prominence: a reappraisal of the foundations of biology. J R Soc Interface 2014; 11:20131017; PMID:24554573; <http://dx.doi.org/10.1098/rsif.2013.1017>
- [185] Ben Jacob E, Becker I, Shapira Y, Levine H. Bacterial linguistic communication and social intelligence. Trends Microbiol 2004; 12:366-72; PMID:15276612; <http://dx.doi.org/10.1016/j.tim.2004.06.006>
- [186] Ben Jacob E, Shapira Y, Tauber AI. Seeking the foundations of cognition in bacteria: from Schrödinger's negative entropy to latent information. Physica A 2006; 359:495-524.
- [187] Ben-Jacob E, Lu M, Schultz D, Onuchic JN. The physics of bacterial decision making. Front Cell Infect Microbiol 2014; 4:154; PMID:25401094; <http://dx.doi.org/10.3389/fcimb.2014.00154>
- [188] Bray D. Wetware: a Computer in Every Living Cell. 2009; New Haven; London: Yale University Press.
- [189] Cox RP, Krauss MR, Balis ME, Dancis J. Studies on cell communication with enucleated human fibroblasts. J Cell Biol 1976; 71:693-703; PMID:993266; <http://dx.doi.org/10.1083/jcb.71.3.693>
- [190] Hämmerling J. Nucleo-cytoplasmic interactions in *Acetabularia* and other cells. Ann Rev Plant Physiol 1963; 14:65-92; <http://dx.doi.org/10.1146/annurev.pp.14.060163.000433>
- [191] Chapman CJ, Nugent NA, Schreiber RW. Nucleic acid synthesis in the chloroplasts of *Acetabularia mediterranea*. Plant Physiol 1966; 41:589-92; PMID:5932403; <http://dx.doi.org/10.1104/pp.41.4.589>
- [192] Nagel T. Mind and Cosmos: Why the Materialist Neo-Darwinian Conception of Nature is Almost Certainly False. Oxford University Press; 2012.
- [193] Searle J. Theory of mind and Darwin's legacy. Proc Natl Acad Sci USA 2013; 110: 10343-8; PMID:23754416; <http://dx.doi.org/10.1073/pnas.1301214110>
- [194] Bastian B, Jetten J, Hornsey M, Leknes S. The positive consequences of pain: a biopsychosocial approach. Pers Soc Psychol Rev 2014; 18:256-279; PMID:24727972; <http://dx.doi.org/10.1177/1088868314527831>
- [195] Heil M. Damaged-self recognition as a general strategy for injury detection. Plant Signal Behav 2012; 7:576-80; PMID:22516811; <http://dx.doi.org/10.4161/psb.19921>
- [196] Heil M, Land WG. Danger signals - damaged-self recognition across the tree of life. Front Plant Sci 2014; 5:578; PMID:25400647; <http://dx.doi.org/10.3389/fpls.2014.00578>
- [197] Heil M, Ibarra-Laclette E, Adame-Álvarez RM, Martínez O, Ramirez-Chávez E, Molina-Torres J, Herrera-Estrella L. How plants sense wounds: damaged-self recognition is based on plant-derived elicitors and induces octadecanoid signaling. PLoS One 2012; 7:e30537; PMID:22347382; <http://dx.doi.org/10.1371/journal.pone.0030537>
- [198] Trewavas AJ, Baluška F. The ubiquity of consciousness. EMBO Rep 2011; 12:1221-5; PMID:22094270; <http://dx.doi.org/10.1038/embor.2011.218>
- [199] Vandekerckhove M, Bulnes LC, Panksepp J. The emergence of primary anoetic consciousness in episodic memory. Front Behav Neurosci 2014; 7:210; PMID:24427125; <http://dx.doi.org/10.3389/fnbeh.2013.00210>
- [200] Fabbro F, Aglioti SM, Bergamasco M, Clarici A, Panksepp J. Evolutionary aspects of self- and world consciousness in vertebrates. Front Hum Neurosci 2015; 9:157; PMID:25859205; <http://dx.doi.org/10.3389/fnhum.2015.00157>